

The hard QCD pomeron: some aspects of its phenomenology and interactions

G.P. Vacca

Department of Physics, University of Bologna and INFN, Italy

Contents

Introduction	4
1 Pomeron and the Regge limit in perturbative QCD	11
1.1 The BFKL equation	13
1.2 Solution for the pomeron amplitude	16
1.3 Conformal Invariance	18
1.4 The colour dipole approach in the large N_c limit	22
2 Hard pomeron with a running coupling constant: a phenomenological approach	27
2.1 Pomeron equation with running coupling	29
2.1.1 Basic equations. Pomeron at $q = 0$	32
2.1.2 Pomeron at $q \neq 0$: the slope	34
2.1.3 Numerical procedure and results	35
2.1.4 Pomerons and the high-energy scattering.	38
2.2 Inclusive jet production.	41
2.2.1 General formalism.	41
2.2.2 Numerical results.	45
2.3 Evolution of the gluon density.	48
2.3.1 Coupling to the virtual photon	49
2.3.2 The initial distribution	52
2.3.3 Numerical results	54
3 Unitarization in the large N_c limit: the three pomeron vertex	59
3.1 General unitarization approach	61
3.2 Large N_c limit for the system of four reggeized gluons	62
3.2.1 Leading order in $1/N_c$	62
3.2.2 Next to leading order in $1/N_c$	65
3.3 Conformal invariance	70
3.4 Double pomeron exchange and triple pomeron interaction	73
3.5 Relation to the dipole approach	76
3.6 Higher order densities in the dipole approach	78

4	Conclusions	85
A		89
A.1	Asymptotic behaviour of the wave functions.	89
A.1.1	The momentum space behaviour	89
A.1.2	The coordinate space behaviour	92
A.2	Two gluons coupling to quark loop for non-zero transfer momentum	94
A.3	Coordinate representation of the non-forward G function	96
A.3.1	Proper part	97
A.3.2	Improper part	98
A.3.3	Infrared stable pieces	100
A.4	Some useful relations in graphical notation.	102

Introduction

Strong interactions continue to be a challenge in theoretical physics. They involve all the phenomena regarding hadrons and actually the best suggestion for a model to describe them is given by Quantum Chromodynamics (QCD), which is a gauge quantum field theory based on a local $SU(3)$ symmetry, which satisfies the general requirements of renormalizability and unitarity of the scattering S -matrix. The elementary degrees of freedom of QCD are quarks and gluons and its complexity has prevented the solution of the theory, i.e. finding the spectrum and correlation functions. On one side interesting work is devoted to analyzing numerically the euclidean version of the theory on a lattice, even if until now the available computational power really restrict the dimension of the lattice involved and thus the significance of obtainable results.

On the other hand analytical efforts are done in two directions. Due to the asymptotic freedom property related to the decrease of the interaction strength with the increase of the energy of the particles involved, perturbation theory in the small coupling regime is applied. A perturbative approach has been also considered for other “small” parameters, such as $1/N_c$ [1, 2, 3] where N_c is the number of colours which defines the special unitary gauge group. Typical non-perturbative methods concern condensates of quarks and gluons and topological objects such as instantons; however we are still far from the understanding the nature of composite bound objects in QCD.

One of these objects seems to be the pomeron, originating from the works in Regge theory, before a consistent field theory approach to strong interactions was established with QCD. The pomeron is related, in a hadronic scattering process, to the rightmost singularity (preferably a pole) of the partial wave amplitude in the complex angular momentum plane. The mathematical structure of the amplitude is commonly associated to the presence of an exchanged object with vacuum quantum numbers which can lead to a power-law-increase of the total cross section with the center of mass energy, in such a case in contrast with the unitarity bounds. In general the process is deeply related to non-perturbative phenomena and the only possibility to understand as much as possible of it with perturbative tools is to consider processes involving the scattering of highly virtual photons or onia-particles where the perturbative analysis in terms of elementary quarks and gluons make sense.

At each order of perturbation theory one finds a leading behaviour proportional to a proper power of the logarithm of the center of mass energy ($\sqrt{s}$). The approach which resums these dominant contributions is called leading log(s) approximation (LLA). One of its principal limitations is related to the fact that the coupling constant is considered fixed. The works of Lipatov et al. [4, 5] were devoted to these calculations for a $SU(N_c)$ gauge theory. According to

previous ideas of Regge theory they use the gluon reggeization property, computed considering the dominant virtual corrections and which can be associated to a bootstrap condition related to unitarity, to construct the BFKL equation which describes the interactions in LLA between two reggeized gluons; the contributions to these interactions may be represented by ladder-type diagrams, resummed in a Bethe-Salpeter-like equation. The vacuum quantum number in the t -channel corresponds to the BFKL hard pomeron. Its singularity is not a pole but a branch-cut in the complex angular momentum plane.

In order to go beyond the fixed coupling case a next-to-leading order (NLO) analysis has been carried on in the last years [6, 7, 8]. A quite complicated generalized BFKL equation was found but still it is not clear if such a model is really applicable to the HERA kinematical region or one may need to go beyond the NLO approximation to give satisfactory predictions and interpretations of the experimental data.

All the theoretical predictions may be tested in various experiments in the kinematical Regge region. In the last years particular interest was devoted to the deep inelastic scattering (DIS) processes at HERA where the small x region is probed. Recent obtained measures [9, 10] have shown a sharp rise of the F_2 proton structure function at low x and this fact may be interpreted as a manifestation of the hard pomeron. Much attention was also given to the observed large rate of rapidity-gap events at small x , another possible signature of the pomeron interaction.

The basic ingredients in the LLA analysis and a brief derivation of the BFKL equation are reported in the first chapter of this thesis. A well known problem related to the consistency of the approach is evidenced by noting that the diffusion-like process described by the BFKL equation may spoil the applicability of perturbation theory, since a huge contribution at large energies may involve small transverse momenta, deeply related to a non-perturbative regime in QCD. The other main problem in the BFKL theory is the violation of unitarity due to the high value of its intercept. The remarkable property of global conformal symmetry of the BFKL equation in the transverse two dimensional space is also shown and this permits one to also have a general explicit solution for the system.

An alternative approach in perturbative QCD to describe the scattering of two onia-particles is given by the colour dipole picture [11, 12], which is based on the computation of the corrections to the light-cone wave functions of onia or virtual photons due to the emission of soft gluons strongly ordered in their scaling variables. This implies a leading logarithmic approach which turns out to be equivalent to the BFKL one. In the large N_c limit it is possible to write an equation for the generating functional of the n -dipole distributions [11] which should be considered in general to have unitarity corrections. A brief introduction to the basic equations of this model is given in the last section of the first chapter, since they will be needed in the analysis performed in the third chapter.

The problems related to the infrared sector one finds in the BFKL theory, which is constructed on a small fixed coupling analysis thus far away from the non-perturbative region which instead can be reached in the diffusion process, has motivated many phenomenological works, beyond the NLO analysis, devoted to the modification of the infrared sector of the BFKL equation.

In the second chapter we try to analyze some properties of a phenomenological model which

have the merit of preserving the gluon reggeization property [13, 14] with the insertion of the running coupling in the generalized BFKL equation. The bootstrap condition is the crucial point since in this way unitarity is satisfied to leading order for the production amplitude with one gluon exchange in the t -channel while in all the other phenomenological attempts it is lost. The second input for the model is the compatibility with the predictions of the renormalization group equations for the gluon density (DGLAP) in the double leading log approximation (DLA) The model depends at least on a mass parameter, related to infrared sector, on which the trajectories of all the singularities depend.

The insertion of the running coupling provides a deep change in the properties of the pomeron equation [15]. A part of the spectrum is found to be discrete. The values are found numerically while the corresponding asymptotic behaviour of the eigenfunction can be derived analytically. The parameter dependence can be fixed, for example, by forcing the value of the slope (computed perturbatively) of the leading singularity. For some reasonable values one finds two supercritical pomeron states (corresponding to negative “energy” for the two reggeized gluons bound states in the Schrödinger-like picture). A rough attempt to estimate some effects of unitarity corrections has been made by considering the multipomeron exchange in the scattering of two virtual photons in the large N_c limit, a case for which a corresponding eikonal formulation can be given. As a result, for the parameter choice considered, one finds that these corrections occur at energies much higher than the ones now-a-days reached. However unitarity requires one to include also the effects of the triple pomeron interaction, as is also shown in the third chapter. Thus unitarity effects can still be significant at lower energies.

In the framework of the standard BFKL approach the process of jet production has been extensively studied showing the importance of mini-jet production and the logarithmic rise of the multiplicity. The fixed coupling analysis requires the introduction of an infrared parameter and the inclusions of some unitarity and coherence effects to restore the correct asymptotic behaviour of the differential cross section for large jet transverse momentum. In the model with running coupling considered [16] one does not have, of course, any problem in the infrared sector and finds, due to the asymptotic behaviour of the pomeron states, a correct tail for the cross section at large jet transverse momenta. The calculations have been carried out on taking into account the two supercritical pomeron states.

In order to obtain a better description of the physical processes at present energies a first improvement consists in considering all the states present in the spectrum of the pomeron equation, and not only the leading contributions of the two supercritical states. This program [17] can be achieved by considering the associated evolution equation in $1/x$ (with x the Bjorken variable) for the pomeron wave function. To this end we are forced to address the problem of the coupling of a virtual photon to the gluon in the running coupling framework, since otherwise there is no link, for example, between the gluon distribution in the proton and its structure function. The bootstrap relation, which concerns only the gluon interaction, is of no help. The idea is to find a corrected dipole density distribution for the photon in order to take into account a minimal running coupling effect. We adopt here a phenomenological approach and guess a possible form of the dipole density from the matching in the DLA of the quark distribution (which one has after coupling the gluons to the dipole) with the one obtained by the DGLAP

evolution. Numerical calculations show a still steep increase of the gluon distribution and the singlet proton structure function, but smaller than the one implied by the standard BFKL theory.

In the second part of the thesis we deal with the unitarity problem from a theoretical point of view. There is a limitation to the validity of the BFKL (fixed coupling) theory for large s or equivalently large $1/x$ and to extend the kinematical region of applicability of the model one has to go beyond the LLA. This problem is shared also by the DGLAP formalism and to solve it one needs to find a systematic way to include unitarity corrections. Continuing the restriction to the onia-particles or virtual photons scattering there is the hope to reach some new insights by means of perturbative tools even if the complete understanding surely requires a true non-perturbative approach (the idea of grasping non-perturbative effects with the help of a perturbative approach can be seen for example in the papers about renormalons).

In order to understand which kind of corrections one has to deal with, it is very useful to consider the evolution of the parton densities in the rapidity. For example it is well known for DIS processes that at lower x the photon probes in the proton a higher density of partons, of typical size $1/Q$ at virtuality Q^2 . The resulting cross section then increases; the unitarity corrections correspond in this picture to the fusion of partons which diminishes the rate of increase of the parton density. A first quantitative model for such a process was given by Gribov, Levin and Ryskin [18] by constructing a modified non linear BFKL equation generating a fan diagrams structure (based on triple ladder vertices). The key point is the presence of multi-parton states in the t -channel and eventually also multi-scattering processes between the partons of the two scattering onia have to be considered as suggested by the dipole picture.

To this end a systematic approach to take into account a minimal set of corrections necessary to restore unitarity has been studied by Bartels [19]. Higher order amplitudes, which contain any number of (reggeized) gluons in the t -channel, are constructed using unitarity and dispersion relations starting from the lower order ones and one has a coupled system of equations for the n reggeized gluon state amplitudes. The main feature is the presence of transitions vertices which change the number of reggeons in the t -channel and thus leads to a reggeon field theory.

These corrections have been studied by considering a system of up to four reggeized gluons in the t -channel [20]. Among the building blocks an effective vertex for the transition between 2 and 4 gluons states in the t -channel appears. In [21] its conformal invariance property was proven and in the appropriate color subspace it is related to the triple pomeron vertex. Another element of some interest is the 4-gluon interacting state (and its generalizations to n gluons) which, as a system with a fixed number of reggeized gluons in the t -channel, can be studied by means of the so called BKP equation [22, 23]. The related amplitude has been studied in the large N_c limit and the system has been proven to be complete integrable and equivalent to an XXX Heisenberg zero spin model [24, 25, 26]. These elements have suggested an interpretation in terms of correlation functions of a 2D conformal field theory (CFT) [27, 28] and one hopes to have an effective formulation in terms of a (2+1) dimensional CFT for QCD in the Regge limit.

In the third chapter some efforts are addressed to obtaining a clearer picture of the interactions present in the system of 4 reggeized gluons in the t -channel in the large N_c limit [29, 30], in particular on having some understanding of the relation between multipomeron exchange and

triple pomeron interactions and making a connections between this reggeon analysis based on the s -channel unitarity and the colour dipole picture for QCD.

We shall study the formal general system of equations for the 4 gluon system at leading and next-to-leading order in $1/N_c$. The double pomeron amplitude at NLO is studied to understand the role of the direct double pomeron exchange (DPE) and the triple pomeron interaction (TPI). The TPI term is analyzed in terms of 3 point function building blocks (a generalization to the non-forward case of a function previously introduced in [20]) to describe in simpler terms the conformal invariance properties of the system. It is later shown in detail the fact, already noted for $N_c = 3$ in [20], that the DPE contribution can be substituted by a completely equivalent TPI term while the inverse is not true and there is a complete agreement with the vertex form in [20] in the limit $N_c \rightarrow \infty$. On coupling to pomerons and working in the coordinate representation one can compare the results obtained from s -channel unitarity with the ones found by R. Peschanski in [31] starting from the colour dipole picture. A substantial agreement is found, apart from an asymmetry factor in the triple pomeron vertex and some single pomeron contributions absent in the dipole approach. A final section is devoted to studying some properties of higher order dipole densities in the A.H. Mueller approach which are found to be represented by a set of pomeron fan diagrams with only a triple pomeron coupling. It is expected that the s -channel unitarity analysis for a larger number of reggeized gluons in the t -channel might tell us the form of higher order pomeron vertices and thus also if the dipole picture has to be improved, as already suggested [31].

The thesis is structured in the following way.

In the first chapter we review some important results of QCD in LLA which are at the basis of the developments in the second and third chapters. In section 1.1 the BFKL equation is briefly derived and in the following one the pomeron solution with some properties presented. Section 1.3 is devoted to show in some details the conformal invariance property of the BFKL equation and finally in section 1.4 a minimal review of the colour dipole approach in the A.H. Mueller formulation is given. In the second chapter we analyze a phenomenological model for considering the running coupling in the hard pomeron. In section 2.1 the definition of the model is given and the properties of the pomeron studied [15]; in particular in section 2.1.1 and 2.1.2 the problem is formulated to make possible a numerical analysis, performed in section 2.1.3, while multi-scattering effects are considered in section 2.1.4. In section 2.2 the inclusive jet production process is considered [16]. Section 2.2.1 is devoted to set up the formalism and to make asymptotic estimates while section 2.2.2 collects some numerical results. The evolution of the gluon density [17] is considered in section 2.3 and in particular in section 2.3.1 we address the problem of the coupling to the virtual photon, in section 2.3.2 we look at the starting distribution and in section 2.3.3 some numerical calculations are shown. Finally in the third chapter the triple pomeron vertex is analyzed in the large N_c limit [30]. In section 3.1 we introduce some basic elements of the unitarization program and in section 3.2 we present the analysis for the 4 gluons system in the large N_c limit, in particular at the leading order in 3.2.1 and at the NLO in 3.2.2 where we establish the form of the DPE and TPI terms. In section 3.3 the properties of conformal invariance are re-analyzed in terms of a simple function and in section 3.4 the interplay between the DPE and the TPI terms is analyzed in detail. In section 3.5

the coupling to pomerons and the relation to the dipole approach is studied while section 3.6 is devoted to study the solutions for general higher order dipole distributions in the standard dipole picture and analyze the triple pomeron interactions structure resulted. In the appendix one can find some more technical parts. In A.1, related to the phenomenological model considered in the third chapter, the asymptotic form of the bound state solution of the pomeron equation is derived. In A.2 the dipole density in the non-forward case for the coupling of two gluons to a quark loop is given. In A.3 the coordinate representation of the function G , needed in chapter 4, is calculated and its infrared stability explicitly shown. In A.4 some relations, used in the fourth chapter, are derived in a graphical notation.

Chapter 1

Pomeron and the Regge limit in perturbative QCD

To understand the concept of Pomeron it is necessary to go back to the old Regge theory and Pomeron field theory. The Pomeron is the rightmost singularity in the complex angular momentum plane of the partial wave signatured amplitude of a hadronic scattering process characterized by the quantum number of the vacuum exchanged in the t -channel. The singularity can be associated to a rise of the cross-section with the center of mass energy, thus leading to a violation of unitarity which can be restored only if more complicated processes than a single pomeron exchange are taken into account. The understanding of the strong interactions in the Regge limit is therefore deeply connected to unitarity from a general point of view.

The study of strong interactions as described by QCD theory is well known to be an incredibly difficult problem and due to the asymptotic freedom which characterizes QCD many efforts have been directed in perturbative analysis, hoping to not lose some important aspects, which could serve as a guideline to understand the features of the complete real QCD.

In particular the concept of the Pomeron has been intensively studied in perturbative QCD in the Regge limit for a long time [4]. At the basis of the approach is the property of Reggeization of the gluon. In general the poles in the complex angular momentum plane of the partial wave amplitude depend on the exchanged t squared momenta so that they may describe a family of exchanged particles of integer even or odd angular momenta, thus rising with spin and mass. An elementary particle, when exchanged in the t channel, gives a non-analytic contribution to the partial amplitude dependence in the angular momentum, and the modern idea of reggeization is related to the fact that summing all the contributions to the amplitude not containing poles in the t -channel leads to a partial wave amplitude with a pole in the t -channel which is located at the spin of the elementary particles if t is taken to be the mass squared of the particle itself. This was verified for the gluon (massive after considering the Higgs mechanism) up to the 8-th order (and partially up to 10-th) in perturbation theory by Cheng and Lo [32], after the work of Grisaru [33] where a first confirmation that the gluon reggeizes was given.

Another method [4] which has been used to check gluon reggeization consists in taking a gluon ladder diagram, constructed with reggeized gluons on the basis of elastic unitarity (the on shell condition turns the reggeized gluons into elementary gluons). On summing these ladder

diagrams one obtains a partial wave amplitude which, in the colour channel, has the same reggeized gluon Regge pole. This is called a “bootstrap” condition and it is at the basis of the BFKL theory and all its extensions.

The analysis of the processes in the Regge limit in perturbation theory reduces to summing the leading log terms. In fact for a small fixed coupling g and high c.m.s. energy s such that $s \rightarrow \infty$, $g \rightarrow 0$ and $g^2 \ln s \sim 1$ the Regge pole contribution has a behaviour $s^{\alpha(t)}$ where $\alpha(t) = 1 + \omega(t)$ is the trajectory and $\omega(t)$ is of second order in the coupling constant g . Therefore a perturbative expansion leads one to consider amplitudes of the order $g^2 s (g^2 \ln s)^n$.

Apart from considering a fixed coupling, its smallness is generally ensured in the Lipatov perturbative approach by giving to the gluon a mass $M \gg \Lambda$ using the Higgs-Brout-Englert mechanism [34, 35]. Thus one has also an infrared regularization but we should remember that in this way one is not really working with true QCD since it is not possible to safely perform the $m \rightarrow 0$ limit without avoiding the non-perturbative region.

The LLA analysis of the amplitude for gluon-gluon scattering in the vacuum channel was started in [4, 5]. Multiparticle production and unitarity were the ingredients used to construct a Bethe-Salpeter equation for ladder contributions to the partial wave amplitude with reggeized gluons in the t -channel. The resulting singularity was found to be a branch cut and not a pole in the complex angular momentum and the BFKL pomeron to be a “bound” state of two reggeized gluons. In the colour octet channel instead the Regge pole ansatz for the gluon applied to the BFKL equation gave directly the self-consistency bootstrap condition.

The BFKL equation can be seen as a Schrödinger-like equation in the transverse space with the evolution flow in the rapidity variable. We just remember, beyond the violation of unitarity, an intrinsic problem in the BFKL perturbative approach due to the diffusion of the gluon momenta in the ladders. For a highly virtual projectile (Q^2) the scattering is characterized by a distribution which tends to be Gaussian in $\ln \mathbf{k}^2$ with an average value $\ln Q^2$ and a width proportional to $\sqrt{\ln s}$, thus with a support which can be strongly present in the non-perturbative region.

The opportunity of applying the results found in this perturbative approach to analyze some real processes is based on the possibility of separating in them the contribution of short distance interactions (hard scales), where perturbation theory can be applied, from the long distance ones (soft scales) where perturbation theory is meaningless.

This is the meaning of the factorization theorems. A first kind used to study many hard processes characterized by a hard scale Q^2 at which the coupling is small is the collinear factorization theorem [36]. An object such as, for example, a structure function in DIS can be written as a convolution between a cross section for an hard parton subprocess (perturbatively calculable and depending on the process virtuality) and a distribution for the parton probed in the hadron involved. The latter distribution depends on the virtuality chosen to probe the hadron by means of a renormalization group equations (DGLAP [37]) so that a non-perturbative input at a lower scale Q_0^2 is necessary to provide the initial conditions for these equations (which are constructed on all the collinear logarithmic contributions). The DGLAP equations can be associated, in the physical gauge, with summing all the contributions of ladder graphs with strongly ordered parton momenta but, in the small x (high energy) limit, higher order contributions, beyond the

leading logs, are found to be significant. This fact has led to consider the perturbative resummation of contributions of the order $(\alpha_s \ln 1/x)^n$ which correspond to the ladder graphs with reggeized gluons in the t -channel and no ordering in the transverse momenta. In this framework a new $\mathbf{k}$ -factorization theorem has been given [38].

We shall introduce in the next section the BFKL equation and show some of its properties and the solutions in the subsequent one. In the third section the proof of its conformal invariance is given, a feature which turns out to be very interesting and also characterizes the symmetry of the three pomeron vertex. We shall return to these properties in the last chapter. The Colour dipole approach in the A. H. Mueller formulation [11, 12] is briefly reviewed in the last section since its use is pertinent in the high energy (and a large number of colours) limit and it has been shown to give an alternative formulation to the hard pomeron phenomena. In chapter 3 we shall try to compare the pomeron interactions described in the two formulations to see the degree of equivalence of the two approaches.

1.1 The BFKL equation

The main results about the LLA are briefly derived in the following [5, 39]. We consider the absorptive part $A_s(s, t) = \text{Im}A(s, t)$ of the dominant non-helicity-flip amplitude for the scattering of two gluons in the limit $s \rightarrow \infty$, t finite and $\alpha_s = g^2/(4\pi) \ll 1$ in the t -channel colour state R . Using unitarity one can compute this quantity by means of the amplitude for the production of $(n+2)$ gluons from two:

$$\text{Im}A^{(R)}(s, t) = \frac{P^{(R)}}{2} \sum_n \int d\tau_{n+2} A^*(n+2, p'_1, p'_2) A(n+2, p_1, p_2) \quad (1.1)$$

where $P^{(R)}$ is the colour projector, $p_1 = (p_+, 0, 0_\perp)$ and $p_2 = (0, p_-, 0_\perp)$ are the momenta of the incoming gluons in the c.m.s. in the light cone representation, so that $s = 2p_1 \cdot p_2 = 2p_+^2$ and $d\tau_{n+2}$ is the $(n+2)$ particle phase measure given by

$$d\tau_{n+2} = \prod_{i=0}^{n+1} (d^3k_i / (2\pi)^3 2k_{i+}) (2\pi)^4 \delta^4(\sum k_i - p_1 - p_2) \quad (1.2)$$

The momenta k_i of the intermediate particles satisfy an on-shell condition. It is convenient to use the Sudakov variables α_i and β_i such that $k_i = \alpha_i p_1 + \beta_i p_2 + k_{i\perp}$, $\alpha_i \beta_i = m_{i\perp}^2/s$ with $m_{i\perp}^2 = m^2 - k_{i\perp}^2$. Other useful variables are the transferred momenta $q_i = p_1 - \sum_{j=0}^{i-1} k_j$ ($k_i = q_i - q_{i+1}$) and the partial energies carried by two adjacent intermediate real particles $s_{i,i+1} = (k_i + k_{i+1})^2$.

In the Regge limit the dominant leading $\log(s)$ contribution comes from the region in the phase space characterized by

$$\alpha_i \gg \alpha_{i+1} \quad , \quad \beta_i \ll \beta_{i+1} \quad (1.3)$$

and thus one has approximatively the equalities $s_{i,i+1} = (\alpha_i/\alpha_{i+1})m_{i+1,\perp}^2$ and $q_i^2 = q_{i+1,\perp}^2$. Changing the variables which represent the measure in phase space to q_i and $s_{i,i+1}$ and performing an

integration in the transverse space and the integrations w.r.t. α_0 and α_{n+1} one finally may write

$$d\tau_{n+2} = \frac{\pi}{s} \prod_{i=1}^{n+1} \frac{d^2 q_i ds_{i-1,i}}{(2\pi)^3} \delta\left(\prod_{i=0}^n s_{i,i+1} - s \prod_{i=1}^n m_{i\perp}^2\right) \quad (1.4)$$

Let us now consider the amplitude for the production of $(n+2)$ gluons. In particular we shall denote by λ_i , a_i and λ'_i , a'_i the helicities and colours of the two incoming gluons and of the two outgoing gluons with nearly the same momenta as the initial ones and by e_i and d_i the polarizations and colours of the remaining n gluons.

In the Regge limit the virtual corrections to the tree diagrams may be analyzed by looking at the one-loop corrections of the scattering process $gg \rightarrow gg$ with both the diagrams related to the s -channel and u -channel physical region. One finds a correction of the type $\alpha(t) \ln s$ and it is natural to make the reggeization ansatz for a correction $(s/s_0)^{\alpha(t)}$ in all orders in α_s in the LLA. The trajectory found for the gluon is $\alpha(t) = 1 + \omega(t)$ with

$$\omega(t) = N_c \alpha_s t \int \frac{d^2 l}{4\pi^2 l^2 (q-l)^2} \quad ; \quad t = -q^2 \quad (1.5)$$

Alternatively one may use a general reggeization ansatz for the gluons in the t -channel and find from the bootstrap condition the trajectory, for a case of the two gluons amplitude in the colour adjoint representation; this confirms the fact that the gluon reggeizes.

The real corrections in the LLA leads one to consider a three gluons reggeon-physical-reggeon vertex (the reggeons in the t -channel) whose explicit expression is

$$F(q_{i+1}, q_i) = -q_{i\perp} - q_{i+1,\perp} + p_1 \left(\frac{k_i p_2}{p_1 p_2} + \frac{q_{i\perp}^2}{k_i p_1} \right) - p_2 \left(\frac{k_i p_1}{p_1 p_2} + \frac{q_{i+1,\perp}^2}{k_i p_2} \right) \quad (1.6)$$

It is orthogonal ($F(q_{i+1}, q_i) k_i = 0$) to the emitted gluon momentum. From the analysis of the two physical-reggeon-physical gluons vertices (external in the ladder) the dominant contribution is helicity conserving and proportional to s so that the amplitude $A_{2 \rightarrow 2+n}$ can be written as

$$A_{2 \rightarrow 2+n} = 2sg\delta_{\lambda_1 \lambda'_1} T_{a'_1 a_1}^{i_1} s_{01}^{\omega(t_1)} t_1^{-1} g T_{i_2 i_1}^{d_1} (e_1 F(q_2, q_1)) s_{12}^{\omega(t_2)} t_2^{-1} g T_{i_3 i_2}^{d_2} (e_2 F(q_3, q_2)) \dots \\ \dots s_{n,n+1}^{\omega(t_{n+1})} t_{n+1}^{-1} g \delta_{\lambda_2 \lambda'_2} T_{a'_2 a_2}^{i_{n+1}} \quad (1.7)$$

where $t_i = -q_i^2$. In order to obtaining an expression for (1.1) it is necessary to sum over the polarizations and the colours of the intermediate real gluons and consider the projection onto the desired colour subspace (we shall be mainly interested in the vacuum state).

The sum over polarizations leads to an effective interaction vertex between the two reggeized gluons. Starting from $\sum_{\lambda_i} e_{\lambda_i}^{\mu_i} e_{\lambda'_i}^{\nu_i} F_{\mu_i}(q_i, q_{i+1}) F_{\nu_i}(q'_i, q'_{i+1})$ and performing the sum over the polarizations (actually using the replacement $\sum_{\lambda_i} e_{\lambda_i}^{\mu_i} e^{\nu_i} \rightarrow -g^{\mu_i \nu_i}$) one obtains

$$F \cdot F' = -2K_q(q_i, q'_i, q_{i+1}, q'_{i+1}) \quad ; \quad K_q(q_i, q'_i, q_{i+1}, q'_{i+1}) = q^2 - (q_{1\perp}^2 q'_{2\perp}^2 + q_{2\perp}^2 q'_{1\perp}^2)/k_{\perp}^2 \quad (1.8)$$

Regarding the colour structure in the t -channel, the gluon-gluon amplitude has to be de-

composed into irreducible representations of the $(N_c^2 - 1) \otimes (N_c^2 - 1)$; by means of a projection operator one has

$$A_{a_2 a'_2, a_1 a'_1}(s, t) = \sum_{(R)} P_{a_2 a'_2, a_1 a'_1}^{(R)} A^{(R)}(s, t) \quad (1.9)$$

For each real gluon emission one has a colour factor which depends on the representation R according to the decomposition of the product of the two corresponding colour matrices

$$\left[\sum_{d_i} T_{b_{i+1}, b_i}^{d_i} (T_{b'_{i+1}, b'_i}^{d_i})^* \right]^{(R)} = \left[-(TT')_{b_{i+1} b'_{i+1}, b_i b'_i} \right]^{(R)} = \lambda_R \delta_{b_{i+1} b_i} \delta_{b'_{i+1} b'_i} \quad (1.10)$$

so that the interaction term is given by

$$\tilde{K}^{(R)} = \lambda_R K_q \quad (1.11)$$

Also for the first and last intermediate gluons one has the same colour factor. In particular one has for the vacuum channel $\lambda_1 \equiv \lambda_0 = N_c$ and for the adjoint representation $\lambda_{N_c^2-1} = N_c/2$.

The absorptive amplitude (1.1) can thus be written in the form

$$\begin{aligned} \text{Im} A^{(R)(n)}(s, t) &= \pi s g^{2(n+2)} \lambda_R^2 \int \prod_{i=1}^{n+1} \frac{d^2 q_i ds_{i-1, i}}{2(2\pi)^3} \left(\frac{s_{i-1, i}}{s_0} \right)^{\omega(t_i) + \omega(t'_i)} \frac{1}{t_i t'_i} \\ &\quad \prod_{i=1}^n 2\tilde{K}_q^{(R)}(q_i, q_{i+1}) \delta \left(\prod_{i=0}^n s_{i, i+1} - s \prod_{i=1}^n |k_{i\perp}^2| \right) \end{aligned} \quad (1.12)$$

and has the signature fixed by the R representation (positive for symmetric and negative for antisymmetric representations). Taking the Mellin transform one obtains for it an asymptotic contribution (which thus corresponds to the Mellin transform nearby the rightmost singularity)

$$\begin{aligned} a_j^{(R)(n)}(t) &= 2g^{2(n+2)} \lambda_R^2 \int \prod_{i=1}^{n+1} \frac{d^2 q_i}{16\pi^3 t_i t'_i (j-1 - \omega(t_i) - \omega(t'_i))} \\ &\quad \prod_{i=1}^n |k_{i\perp}^2|^{j-1} 2K_q^{(R)}(q_i, q_{i+1}) \end{aligned} \quad (1.13)$$

We note that in the validity region (j close to 1) the $|k_{i\perp}^2|^{j-1}$ factor may be omitted. Changing the notation to the 2-dimensional transverse vectors $l_{1i} = q_{i\perp}$, $l_{2i} = -q'_{i\perp}$ such that $l_{1i} + l_{2i} = q$, $t_i = -l_i^2$ and $t = -q^2$ one may write in an operatorial form

$$a_j^{(R)(n)}(t) = 2g^2 \lambda_R^2 \int \frac{d^2 l_{1,1}}{16\pi^3 t_{11} t_{21}} \langle l_{11} | D(V^{(R)} D)^n | l_{1, n+1} \rangle \frac{d^2 l_{1, n+1}}{16\pi^3} \quad (1.14)$$

where we have defined the operators

$$D(l_1) = \frac{1}{j-1 - \omega(t_1) - \omega(t_2)} \quad ; \quad V^{(R)}(l_1, l'_1) = 2 \frac{\tilde{K}_q^{(R)}(l_1, l'_1)}{l_1^2 l_2'^2} \quad (1.15)$$

It is now very easy to sum the contributions of all the ladder diagrams and one has

$$a_j^{(R)}(t) = 2g^2\lambda_R^2 \int \frac{d^2l_{1,1}}{16\pi^3} \frac{\phi_{jq}^{(R)}}{t_{11}t_{21}} \quad (1.16)$$

with

$$\phi_{jq}^{(R)}(l_1) = \int \frac{d^2l'_1}{16\pi^3} \langle l_1 | D(1 - V^{(R)}D)^{-1} | l'_1 \rangle \quad (1.17)$$

The Lipatov equation is a Bethe-Salpeter equation for $\phi_{jq}^{(R)}(l_1)$ which can be obtained by taking the matrix element of the operatorial identity $D^{-1}T^{(R)} = 1 + V^{(R)}T^{(R)}$ w.r.t. the states $\langle l_1 |$ and $|l'_1\rangle$ where $T^{(R)} = D(1 - V^{(R)}D)^{-1}$ and integrating w.r.t. l'_1 . It reads

$$[j - 1 - \omega(t_1) - \omega(t_2)]\phi_{jq}^{(R)}(l_1) = 1 + 2\alpha_s\lambda_R \int \frac{d^2l'_1}{4\pi^2} V_q(l_1, l'_1)\phi_{jq}^{(R)}(l'_1) \quad (1.18)$$

with the interaction potential given by

$$V_q(l_1, l'_1) = \left(\frac{l_1^2}{l'^2_1} + \frac{l_2^2}{l'^2_2}\right) \frac{1}{(l_1 - l'_1)^2} - \frac{q^2}{l'^2_1 l'^2_2} \quad (1.19)$$

This equation in the gluon channel gives the bootstrap condition if one requires that the solution of this equation must have a Regge pole of a reggeized gluon, i.e. at $j = 1 + \omega(t)$. Denoting by $c_q(l_1)$ the residue of the solution at this pole, the bootstrap condition can be written for $N_c = 3$ as

$$[\omega(t) - \omega(t_1) - \omega(t_2)]c_q(l_1) = 3\alpha_s \int \frac{d^2l'_1}{4\pi^2} V_q(l_1, l'_1)c_q(l'_1) \quad (1.20)$$

The gluon trajectory found by perturbative calculations corresponds to a residue c_q independent of l_1 . Hence in the gluon channel the whole solution for $\phi_{jq}^{(8_A)}(l_1)$ is determined by this pole.

1.2 Solution for the pomeron amplitude

The Lipatov equation can be written as an inhomogeneous Schrödinger equation $(H - E)\phi = f$ with an “energy” $E = 1 - j$ and an Hamiltonian

$$H = -\omega(l_1) - \omega(l_2) - 2\alpha_s\lambda_R V_q \quad (1.21)$$

In the vacuum case one has $\lambda_R = N_c$ and one may check that for $q \neq 0$, as will be shown also in the next section, the infrared divergences cancels between the gluon trajectories and the interaction potential term. For $q = 0$ the divergences at $l'_1 = l'_2 = 0$ remain but actually disappear for the coupling of the pomeron to a colourless physical object, since the function $\phi_{jq}(l \rightarrow 0, q)$ vanishes.

Let us consider the forward scattering case ($q = 0$). It is convenient to consider a new

function $\psi(l_1) = \phi(l_1)/l_1^2$ for which one can write the equation

$$[j - 1 - 2\omega(t_1)]\psi(l_1) = \frac{1}{l_1^2} + 4\alpha_s\lambda_R \int \frac{d^2l'_1}{4\pi^2} \frac{\psi(l'_1)}{(l_1 - l'_1)^2} \quad (1.22)$$

The interaction term is of the Coulomb type and both the trajectory and the potential can be studied in the limit of zero regularizing mass. Using the expressions given in the next section (1.35) and (1.37) one may write the Hamiltonian for the vacuum colour channel ($\lambda_0 = N_c = 3$)

$$H = (3\alpha_s/\pi)h \quad ; \quad h = \ln l^2 + \ln \rho^2 - 2(\ln 2 + \psi(1)) \quad ; \quad E = 1 - j = (3\alpha_s/\pi)\epsilon \quad (1.23)$$

where the dependence on the gluon mass has disappeared and $\psi(1)$ is referred to the logarithmic derivative of the Γ function. To find the eigenvalues and eigenfunction of this Hamiltonian one can take advantage of the scaling symmetry $\rho \rightarrow c\rho$, $l \rightarrow l/c$ which is manifest. The symmetry between the coordinate and the momentum space is complete and one may write easily the proper functions for the system both in coordinate and momentum space

$$\psi_{\nu n}(\rho) = a_{\nu n} \rho^{-1+i\nu} e^{in\phi} \quad ; \quad \psi_{\nu n}(l) = b_{\nu n} l^{-1-i\nu} e^{in\phi} \quad (1.24)$$

for any real ν and integer n . It is also easy to show that these functions are orthonormalized if $|a_{\nu n}|^2 = |b_{\nu n}|^2 = 1/4\pi^2$. Defining $\mu = (1/2)(1 + n + i\nu)$ the relation between the two normalizing factors is for $n \geq 0$

$$b_{\nu n} = a_{\nu n} i^n 2^{-i\nu} \frac{\Gamma(\mu^*)}{\Gamma(\mu)} \quad (1.25)$$

while the $n < 0$ case is obtained by complex conjugation. The eigenvalues are computed through the definition $h\psi_{\nu n} = \epsilon_{\nu n}\psi_{\nu n}$; in particular one has

$$\epsilon_{\nu n} = 2\text{Re} \psi\left(\frac{1 + |n| + i\nu}{2}\right) - 2\psi(1) \quad (1.26)$$

The minimal value $\epsilon_0 = -4 \ln 2$ is reached at $\nu = 0$ and $n = 0$ and gives for the BFKL pomeron the intercept

$$j = 1 + \frac{3\alpha_s}{\pi} 4 \ln 2 \quad (1.27)$$

From this value the violation of the Froissart bound [41, 42, 43] derived from unitarity is evident. The pomeron Green function in the forward direction can be easily constructed considering the inhomogeneous term of the equation for ψ , $f = (2\pi)^2 \delta(l - l') l^{-2} l'^{-2}$, and in momentum space is

$$G_\epsilon(l, l') = \frac{1}{4\pi^2 l^3 l'^3} \sum_{n=-\infty}^{\infty} e^{in(\phi - \phi')} \int d\nu \left(\frac{l'}{l}\right)^{i\nu} \frac{1}{\epsilon_{\nu n} - \epsilon} \quad (1.28)$$

To see which behaviour the solution has in the high momentum (short range) limit one should study the singularities of the Green function in the ν plane. Since we are interested in the dominant contribution, which one has for ϵ close to ϵ_0 , we consider the expansion up to the second order

$$\epsilon_{\nu 0} = \epsilon_0 + a\nu^2 \quad (1.29)$$

where $a = (7/2)\zeta(3)$. Defining $w = \epsilon_0 - \epsilon = (\pi/3\alpha_s)(j-1) - |\epsilon_0|$ the singularity is at $w = 0$ and the asymptotic behaviour for large l is calculated by taking the residue at $\nu = i\sqrt{w/a}$. Thus one obtains

$$\psi_\epsilon(l) = \sum_{n=-\infty}^{\infty} e^{in\phi} \int_{-\infty}^{+\infty} d\nu l^{-1+i\nu} b_{\nu n} \frac{\langle \psi_{\nu n} | f \rangle}{\epsilon_{\nu n} - \epsilon} \simeq \frac{\langle \psi_{00} | f \rangle}{2l\sqrt{aw}} e^{-\sqrt{\frac{w}{a}} \ln l} \quad (1.30)$$

and it is evident that the pomeron singularity is branch cut of type $1/\sqrt{j-1-|\epsilon_0|}$. The leading behavior at high energies s can be found by performing an inverse Mellin transform for the positive signatred amplitude

$$\psi(s, l) = \frac{i}{4} \int dj \psi_j(l) s^j \frac{e^{-i\pi j} + 1}{\sin \pi j} \quad (1.31)$$

giving the dominant contribution

$$\psi(s, l) \simeq \frac{i\pi}{4l} s^{1+\frac{3\alpha_s}{\pi}|\epsilon_0|} \frac{3\alpha_s}{\pi} \frac{\langle \psi_{00} | f \rangle}{\sqrt{3\alpha_s a \ln s}} e^{-\pi \frac{\ln^2 l}{12\alpha_s a \ln s}} \quad (1.32)$$

Thus the average value of l increases with s since the distribution is Gaussian in $\ln l$ with a width proportional to $\sqrt{\ln s}$. This fact is characteristic of a diffusion process in the transverse momenta; the Lipatov equation can in fact be seen as an diffusion equation in the rapidity flow. From this fact we address an important problem in the BFKL theory. Such a diffusion process may spoil the applicability of perturbation theory since at sufficient large energy the distribution in the transverse momenta can have support in the small values region corresponding to large distances and typically regarding a non-perturbative regime.

1.3 Conformal Invariance

The BFKL equation manifests two important properties: the holomorphic separability and conformal invariance. An elegant way to prove them [40, 27, 39] can be achieved by rewriting the Hamiltonian of the two interacting reggeized gluons in the complex coordinate space representation for the transverse plane. In the following we will fix the notation: given two vectors $\mathbf{a} = (a_x, a_y)$ and $\mathbf{b} = (b_x, b_y)$ we have

$$a = a_x + ia_y; \quad , \mathbf{a} \cdot \mathbf{b} = \frac{1}{2}(ab^* + a^*b) \quad (1.33)$$

The complex vectors for the impact parameter r_i will be conjugate to the momenta $l_i = -i\frac{\partial}{\partial r_i}$. It is also convenient to introduce the relative momentum $l = \frac{1}{2}(l_1 - l_2)$ and relative distance $\rho = \mathbf{r}_1 - \mathbf{r}_2$ (vector) and $\rho = \frac{1}{2}(r_1^* - r_2^*)$ (complex number) so that ρ, l and ρ^*, l^* forms, each one, an independent pair of canonical variables and $|\rho|^2 = 4\rho\rho^*$. It should be clear from the context which is, each time, used. Further if in the configuration space we are referring to a vector or to a complex number should be clear from the context. Moreover we define $k = l_1 - l'_1$.

The interaction potential (1.19) can be rewritten as

$$V_q(l_1, l'_1) = l_1 l_2^* \frac{1}{k k^*} (l'_1 l'^*_2)^{-1} + c.c. = l_1 l_2^* W(\rho) (l'_1 l'^*_2)^{-1} + c.c. \quad (1.34)$$

where $W(\rho)$ is a Coulombian potential in the coordinate representation. The explicit form of W , after having introduced an infrared regulator m , is

$$W(\rho) = \frac{K_0(m\rho)}{2\pi} \underset{m \rightarrow 0}{\approx} \frac{1}{2\pi} \left[\psi(1) - \ln \frac{m\rho}{2} \right] \quad (1.35)$$

and consequently one obtains

$$V_q = -\frac{1}{4\pi} \left[l_1 \ln \rho l_1^{-1} + l_2 \ln \rho l_2^{-1} + \ln m^2 - 2\psi(1) \right] + c.c. \quad (1.36)$$

The regularized form of the gluon Regge trajectory reads

$$\omega(l_1) \underset{m \rightarrow 0}{\approx} -\frac{N_c \alpha_s}{2\pi} (\ln l_1 - \ln m + c.c.) \quad (1.37)$$

so that the Hamiltonian is holomorphic separable, i.e. $H = \tilde{H}(\rho, l_i) + c.c.$, and, for the R colour channel,

$$\tilde{H}^R(\rho, l_i) = \frac{N_c \alpha_s}{2\pi} \left[\ln l_1 + \ln l_2 + \frac{\lambda_R}{N_c} (l_1 \ln \rho l_1^{-1} + l_2 \ln \rho l_2^{-1}) + \left(\frac{\lambda_R}{N_c} - 1 \right) \ln m^2 - 2 \frac{\lambda_R}{N_c} \psi(1) \right] \quad (1.38)$$

In the vacuum channel the infrared divergences (regularized by m) in the interaction and the trajectory terms cancel and on rescaling the wave function ϕ in the form $\phi = l_1^2 l_2^2 \psi$, the new Hamiltonian for ψ will be given by $H' = \frac{N_c \alpha_s}{2\pi} (h_{12} + \bar{h}_{12})$ with

$$h_{12} = \ln l_1 + \ln l_2 + l_1^{-1} \ln \rho l_1 + l_2^{-1} \ln \rho l_2 - 2\psi(1) \quad (1.39)$$

The holomorphic separability of the Hamiltonian is an important property and as a consequence the eigenvalues will be given by the sum of two contributions, one the complex conjugate of the other, and the corresponding eigenfunctions will be factorized in two parts, the first depending only on r_1, r_2 and the second part on the conjugate quantities.

Let us now consider a global conformal transformation in two dimension, represented by the Möbius mapping

$$r \rightarrow r' = \frac{ar + b}{cr + d} \quad (1.40)$$

where a, b, c, d are complex number such that $ad - bc = 1$. The generators of the Möbius group are $M^0 = rl$, $M^- = l$, $M^+ = r^2 l$ where $l = -i \frac{\partial}{\partial r}$; M_0 generates the rotation and scale transformations and $M^\pm$ the translations. The Casimir operator is given by $M^2 = M^{0^2} - \frac{1}{2} \{M^+, M^-\}$. For a system of n gluons the generators are given by the direct sum of the generators acting on the different gluon spaces.

The Hamiltonian, which describes the two gluon system, is evidently invariant under the action of the M_0 and $M^\pm$ generators (also h_{12} given in (1.39) is invariant). To prove the

conformal invariance it is convenient [40, 27, 39] to consider the finite transformation defined by $r \rightarrow -\frac{1}{r}$ which leads to

$$\rho \rightarrow \frac{\rho}{r_1 r_2} \quad ; \quad l_i \rightarrow r_i^2 l_i \quad (1.41)$$

and rewrite the Hamiltonian in a different form. Let us therefore consider the following operator identities

$$\begin{aligned} \partial_1 &= \Gamma^{-1}(1+z)\rho^{-1}\Gamma(1+z) & \rightarrow & \ln \partial_1 = -\Gamma^{-1}(1+z)\ln \rho \Gamma(1+z) \\ \rho^2 \partial_1 &= \Gamma(z)\rho\Gamma^{-1}(z) & \rightarrow & \ln \rho^2 \partial_1 = \Gamma(z)\ln \rho \Gamma^{-1}(z) \end{aligned} \quad (1.42)$$

where $\partial_1 = \frac{\partial}{\partial r_1}$ and $z = \rho \partial_1$. They are similarity transformations between ρ^{-1} and ∂_1 and between ρ and $\rho^2 \partial_1$. By noting that $[z, \ln \rho] = 1$ one can use the representation $\ln \rho = -\frac{\partial}{\partial z}$ and derive from the relations (1.42)

$$\begin{aligned} \ln \partial_1 &= -\ln \rho + \psi(z) + \frac{1}{z} \\ \ln \rho^2 \partial_1 &= \ln \rho + \psi(z) \end{aligned} \quad (1.43)$$

Introducing these relations in (1.39) and also substituting the operator identity $l_1^{-1} \ln \rho l_1 = \ln \rho - \frac{1}{z}$ the holomorphic part h_{12} can be written as

$$h_{12} = \ln \rho^2 \partial_1 + \ln \rho^2 \partial_2 - 2 \ln \rho - 2\psi(1) \quad (1.44)$$

In this last expression we made use of the cancellation of imaginary terms between the holomorphic and antiholomorphic part.

It is now easy to prove the invariance of the Hamiltonian with respect to the transformation $r_i \rightarrow -\frac{1}{r_i}$, $i = 1, 2$. In fact applying the transformations (1.41) one has

$$\rho^2 \partial_1 \rightarrow \frac{\rho^2 \partial_1}{r_2^2} \quad ; \quad \rho^2 \partial_2 \rightarrow \frac{\rho^2 \partial_2}{r_1^2} \quad (1.45)$$

and the conformal invariance is verified by direct substitution.

Once the conformal invariance is known to be a property of the system, one can take advantage of it by expanding the general solution of the BFKL equation in terms of the conformal basis functions, which are eigenfunctions of the Casimir operator of the Möbius group and are then related to its irreducible representations.

For a system of two gluon the Casimir operator for the holomorphic part is given by $M^2 = -\rho l_1 l_2$. Using the notation $\rho = r_{12} = r_1 - r_2$ the conformal basis is defined by

$$M^2 \bar{M}^2 E_\mu(r_1, r_2) = \frac{1}{16} r_{12}^4 \partial_1^2 \partial_2^2 E_\mu(r_1, r_2) = \frac{1}{16} \mu(\mu-1) \bar{\mu}(\bar{\mu}-1) E_\mu(r_1, r_2) \quad (1.46)$$

where $\mu = \frac{1-n}{2} + i\nu$ and $\bar{\mu} = \frac{1+n}{2} + i\nu$ are the conformal weights which label the representation; n , the conformal spin, is an integer and ν is a real number. The basis functions (in complex

notation) are explicitly given by

$$E_\mu(r_1, r_2) = E_{n,\nu,r_0}(r_1, r_2) = \left(\frac{r_{12}}{r_{10}r_{20}}\right)^{\frac{1-n}{2}+i\nu} \left(\frac{r_{12}^*}{r_{10}^*r_{20}^*}\right)^{\frac{1+n}{2}+i\nu} \quad (1.47)$$

One notes the presence of an additional quantum number, the coordinate r_0 , in some way related to translational invariance. The corresponding eigenvalues of the Casimir operator can be written conveniently to obtain

$$M^2 \bar{M}^2 E_\mu(r_1, r_2) = \frac{\pi^8}{4a_{n+1,\nu}a_{n-1,\nu}} E_\mu(r_1, r_2) \quad (1.48)$$

where we have used the standard notation

$$a_{n,\nu} \equiv a_\mu = \frac{\pi^4}{2} \frac{1}{\nu^2 + n^2/4} \quad (1.49)$$

The conformal basis forms a complete system:

$$r_{12}^4 \delta^2(r_{11'}) \delta^2(r_{22'}) = \sum_\mu E_\mu(r_1, r_2) E_\mu^*(r'_1, r'_2) \quad (1.50)$$

The notation used regarding the sum over the quantum numbers is the following

$$\sum_\mu = \sum_{n=-\infty}^{\infty} \int d\nu \frac{1}{2a_{n,\nu}} \int d^2 r_0 \quad (1.51)$$

Another important property is the orthogonality relation which contains an additional term due to the equivalence of the representations which correspond to the (n, ν) and $(-n, -\nu)$ quantum numbers. It is given by

$$\begin{aligned} \int \frac{d^2 r_1 d^2 r_2}{r_{12}^4} E_{n,\nu,r_0}(r_1, r_2) E_{n',\nu',r'_0}^*(r_1, r_2) = \\ a_{n,\nu} \delta_{nn'} \delta(\nu - \nu') \delta^2(r_{00'}) + b_{n,\nu} \delta_{n,-n'} \delta(\nu + \nu') |r_{00'}|^{-2-4i\nu} \left(\frac{r_{00'}}{r_{00'}^*}\right)^n \end{aligned} \quad (1.52)$$

where

$$b_{n,\nu} = \pi^3 \frac{2^{4i\nu}}{-i\nu + |n|/2} \frac{\Gamma(-i\nu + (1 + |n|)/2)}{\Gamma(i\nu + (1 + |n|)/2)} \frac{\Gamma(i\nu + |n|/2)}{\Gamma(-i\nu + |n|/2)} \quad (1.53)$$

The proof of these relations can be found in [40]. The spectrum of the BFKL Hamiltonian for the general non-forward case ($q \neq 0$) is the same as the forward case, as a consequence of the global conformal invariance:

$$\omega_\mu = \omega_{n,\nu} = \frac{N_c \alpha_s}{\pi} \left[2\psi(1) - \psi\left(\frac{1 + |n|}{2} + i\nu\right) - \psi\left(\frac{1 + |n|}{2} - i\nu\right) \right] \quad (1.54)$$

The solution of the BFKL equation with an inhomogeneous term Ψ_0 can then be expanded in

terms of the conformal basis functions: in the Mellin representation one has

$$\Psi_j(r_1, r_2) = \sum_{\mu} \frac{E_{\mu}(r_1, r_2) \langle \mu | \Psi_0 \rangle}{j - 1 - \omega_{\mu}} \quad (1.55)$$

and moving to the rapidity space representations, where the BFKL equation is an evolution equation in the rapidity variable, one finds

$$\Psi(r_1, r_2; y) = \sum_{\mu} e^{y\omega_{\mu}} E_{\mu}(r_1, r_2) \langle \mu | \Psi_0 \rangle \quad (1.56)$$

Here we have defined

$$\langle \mu | \Psi_0 \rangle = \int \frac{d^2 r_1 d^2 r_2}{r_{12}^4} E_{\mu}^*(r_1, r_2) \Psi_0(r_1, r_2) \quad (1.57)$$

1.4 The colour dipole approach in the large N_c limit

The light-cone wave function of colourless objects, such as onia or virtual photons, has been studied by many authors [11, 12, 44] to compute the corrections to the wave function given by the emission of soft gluons with scaling variables strongly ordered. The kinematical region $z_{k+1} \gg z_k$ thus relates the results found to the one obtained in the BFKL approach. Incidentally the derivation is quite different and simpler and avoids some suspicious assumptions taken in the construction of the BFKL theory.

The colour dipole approach has also the merit of giving, without further effort, some insight in the multipomeron exchange and interactions as we shall see in chapter 3. In this section we shall sketch a brief derivation of the model in the approach given by A.H. Mueller [11, 12, 39] and show its relation to the BFKL pomeron.

The first object introduced is the wave function of the colourless object (onium or virtual photon) with momentum $(p_+, 0, \mathbf{0})$ in the lowest order in the coupling, with no soft gluons emitted, so that we may think of it as split in a quark-antiquark pair. Let k_1 and $-k_1$ denote the transverse momenta of the antiquark and of the quark respectively; the scaling variable for the antiquark is defined by $z_1 = k_{1+}/p_+$. We shall mainly use the coordinate representation in which the wave function reads

$$\psi_{\alpha\beta}^{(0)}(r_1, z_1) = \int \frac{d^2 k_1}{(2\pi)^2} \psi_{\alpha\beta}^{(0)}(k_1, z_1) e^{ik_1 r_1} \quad (1.58)$$

where α and β are indices for degrees of freedom such as spin and colour. The incoherent sum over spin and colour of the square moduli of the wave function is

$$\Phi^{(0)}(r_1, z_1) = \sum_{\alpha\beta} |\psi_{\alpha\beta}^{(0)}(r_1, z_1)|^2 \quad (1.59)$$

and the normalization will be given by

$$N = \int_0^1 \frac{dz_1}{2z_1(1-z_1)} \int \frac{d^2 r_1}{2\pi} \Phi^{(0)}(r_1, z_1) \quad (1.60)$$

For a hadron N is finite and can be chosen to be 1 while for a photon it would diverge.

Let us consider the contribution to order α_s due to the emission of one soft gluon. One will search for a component of the wave function of the form $\psi_{\alpha\beta}^{(1)a}(r_1, z_1; r_2, z_2)$ where r_2, z_2 and a are the transverse coordinate, the scaling and the colour variables of the soft gluon ($z_2 \ll z_1, 1 - z_1$) respectively. Summing the contributions due to the emission of a soft gluon from the quark and the antiquark one finds

$$\psi_{\alpha\beta}^{(1)a}(r_1, z_1; r_2, z_2) = -i(g/\pi)t^a\psi_{\alpha\beta}^{(0)}(r_1, z_1)[(r_2e)/r_2^2 - (r_{21}e)/r_{21}^2] \quad (1.61)$$

where e is the polarization vector of the soft gluon and $r_{ij} = r_i - r_j$. Introducing the quark coordinate $r_0 = 0$, taking the square modulus of the wave function and the sum over colour and spin one finds the distribution

$$\Phi^{(1)}(r_1, z_1; r_2, z_2) = \Phi^{(0)}(r_1, z_1)c\frac{r_{10}^2}{r_{20}^2r_{21}^2} \quad (1.62)$$

where $c = 4\alpha_s C_F/\pi$, $C_F = (N_c^2 - 1)/(2N_c)$.

In order to consider the contributions due to the multiple emissions of soft gluons some simplifications have been taken into account so that it is possible to find an integral equation which resums all the orders. Precisely the large N_c limit has been considered, under which the gluon lines can be represented in terms of $q\bar{q}$ pairs and only diagrams with a planar topology survive.

For a second soft gluon emitted (characterized by r_3, z_3) one finally obtains

$$\begin{aligned} \Phi^{(2)}(r_1, z_1; r_2, z_2; r_3, z_3) &= \Phi^{(1)}(r_1, z_1; r_2, z_2)c\left(\frac{r_{02}^2}{r_{03}^2r_{32}^2} + \frac{r_{12}^2}{r_{13}^2r_{32}^2}\right) \\ &= \Phi^{(0)}(r_1, z_1)c^2r_{01}^2\left(\frac{1}{r_{02}^2r_{23}^2r_{31}^2} + \frac{1}{r_{03}^2r_{32}^2r_{21}^2}\right) \end{aligned} \quad (1.63)$$

and the formula for an emission of n soft gluons in the kinematical region $z_{n+1} \ll z_n \ll \dots z_2$ is easily inferred

$$\Phi^{(n)}(r_1, z_1; r_2, z_2; \dots r_{n+1}, z_{n+1}) = \Phi^{(0)}(r_1, z_1)c^n r_{01}^2 \left(\frac{1}{r_{12}^2 r_{23}^2 \dots r_{n+1,0}^2} + \text{perm. of } 2, 3, \dots, n+1 \right) \quad (1.64)$$

In the strongly ordered region the integration in the scaling variables of the emitted soft gluons can evidently be done and in the leading order one obtains a trivial logarithmical factor $(\ln z_{10})^n/n!$ where $z_{10} = z_1/z_0$. It is possible to introduce [11] a generating functional for the distribution of the transverse coordinates of the emitted gluons:

$$\Phi^{(n)}(r_1, z_1; r_2, \dots r_{n+1}; z_0) = \Phi^{(0)}(r_1, z_1)(1/n!) \prod_2^{n+1} \frac{\delta}{\delta u(r_i)} Z(r_1, r_0, z_{10})_{u=0} \quad (1.65)$$

The generating functional satisfies the following nonlinear equation

$$Z(r_1, r_0, z_{10}) = 1 + cr_{10}^2 \int_{z_0}^{z_1} \frac{dz_2}{z_2} \int \frac{d^2 r_2}{4\pi r_{02}^2 r_{21}^2} u(r_2) Z(r_2, r_0, z_{20}) Z(r_2, r_1, z_{20}) \quad (1.66)$$

One may be interested in a functional generating not the distribution in terms of the transverse positions of the emitted gluons but instead in terms of r_{ik} , already present in (1.66), which correspond to the dimensions of the colour dipoles. The proper generating functional in this case is

$$D(r_{10}, z_{10}; u) = u(r_{10}) + cr_{10}^2 \int_{z_0}^{z_1} \frac{dz_2}{z_2} \int \frac{d^2 r_2}{4\pi r_{12}^2 r_{20}^2} D(r_{12}, z_{20}; u) D(r_{21}, z_{20}; u) \quad (1.67)$$

The generating functionals given above are very useful because using them one can easily calculate the multiple inclusive distributions for gluons or dipoles. In fact after taking the functional derivatives with respect to u one has just to set its value to 1 instead of 0 as in (1.65). On setting $u = 1$ without taking any derivative one has the probability to find any number of gluons or dipoles emitted with a scaling variable z in the range $z_0 < z < z_1$. These quantities for both (1.66) and (1.67) are evidently not normalized and divergent, the divergence due to the integration in the region of small inter-gluon distances in the transverse plane ($r_{ij} = 0$).

In order to cure these ultraviolet divergences one has to include also the virtual contributions given by loop diagrams; in fact the previous computation was performed only at the tree level. One can even compute the contribution of the virtual corrections by imposing that the divergent part of real and virtual gluon emissions compensate each other to give a normalized probability and checking that the contribution is correct at the lowest loop order. This procedure has been followed for example in [11] and leads to the following equations for the functionals Z and D :

$$Z(r_1, r_0, z_{10}; u) = z_{10}^{2\omega(r_{10})} + cr_{10}^2 \int_{z_0}^{z_1} \frac{dz_2}{z_2} \int \frac{d^2 r_2}{4\pi r_{02}^2 r_{21}^2} u(r_2) z_{12}^{2\omega(r_{10})} Z(r_2, r_0, z_{20}; u) Z(r_2, r_1, z_{20}; u) \quad (1.68)$$

$$D(r_{10}, z_{10}; u) = u(r_{10}) z_{10}^{2\omega(r_{10})} + cr_{10}^2 \int_{z_0}^{z_1} \frac{dz_2}{z_2} \int \frac{d^2 r_2}{4\pi r_{12}^2 r_{20}^2} z_{12}^{2\omega(r_{10})} D(r_{12}, z_{20}; u) D(r_{20}, z_{20}; u) \quad (1.69)$$

where the quantity $\omega(r)$ has the same functional form of the gluon Regge trajectory respect to its momentum argument, i.e.

$$\omega(r_{01}) = -(c/2)r_{01}^2 \int \frac{d^2 r_2}{4\pi r_{02}^2 r_{21}^2} = (2\alpha_s C_F / \pi) r_{01}^2 \int \frac{d^2 r_2}{4\pi r_{02}^2 r_{21}^2} \quad (1.70)$$

The virtual corrections to the functional equations are related to the property of gluon reggeization of the BFKL approach.

Let us now look at the connection between the colour dipole model and the BFKL pomeron. One first obtains an equation for the inclusive one dipole distribution taking a functional derivative with respect to u of relation (1.69) and setting $u = 1$. The resulting equation in term of the

rapidity $y = \ln z_{10}$ reads

$$n(r_{10}, r, y) = e^{2y\omega(r_{10})} \delta^2(r - r_{10}) + 2cr_{10}^2 \int_0^y dy' e^{2(y-y')\omega(r_{10})} \int \frac{d^2 r_2}{4\pi r_{12}^2 r_{20}^2} n(r_{21}, r, y') \quad (1.71)$$

and taking a derivative with respect to y one has the evolution equation

$$\frac{\partial}{\partial y} n(r_{10}, r, y) = 2\omega(r_{10}) n(r_{10}, r, y) + \frac{2\alpha_s C_F}{\pi^2} \int d^2 r_2 \frac{r_{10}^2}{r_{12}^2 r_{20}^2} n(r_{21}, r, y) \quad (1.72)$$

which describes a dual BFKL diffusion in the transverse coordinate space instead of in the transverse momentum space. This equation is also infrared and ultraviolet finite but one may need to regularize it with an ultraviolet cutoff; in it the role of the coordinates and momenta is interchanged.

Chapter 2

Hard pomeron with a running coupling constant: a phenomenological approach

The basic elements in the BFKL approach for the multi-Regge kinematics of Yang-Mills theories are the effective vertices (real contribution) and the Regge trajectory (virtual contribution) whose separate bad behaviour in the infrared sector cancels in the total cross section. In the LLA approximation all the contributions, which are present in the BFKL equation for the pomeron, are based on calculations in the first non-trivial order in perturbation theory. The effective QCD coupling constant (taken to be small since $\alpha_s \ln s \sim 1$) is considered fixed and one does not know the region of applicability of LLA and in particular the energies and momenta fixing the scale for α_s . This is at the basis of the weakness of the LLA approximation, since the quantitative results can be strongly modified by changing the scale of virtuality.

For this reason a program to compute the next-to-leading order corrections to the BFKL kernel of the Lipatov equation has been carried out in the last years [6] with the hope to study soon the so called quasi-Regge kinematics for gluon production. The one loop corrections to the effective vertices, the two loop corrections to the gluon trajectory and terms for the productions of two gluons and quark-antiquark with fixed invariant mass have been computed [7]. The nice feature is that also at the NLO order the pomeron can be seen as a compound state of two reggeized gluons. The generalized BFKL equation however is quite complicated [8] and there is no hope to obtain an explicit analytical solution but the study of such a kernel should be very useful to understand by means of numerical calculations the validity of the predictions in the Regge kinematics and corrections in the quasi-Regge limit. Moreover it is still not clear if such results are applicable in the HERA kinematical region or one should go even beyond the NLO analysis [8].

A part of this well established program some attempts have been made in order to take into account the running of the QCD coupling with a minimal added work. The first idea [27] has been to introduce directly in the BFKL equation for the pomeron the running coupling with some assumptions about the momentum and distance scales at which the gluons interact. Also in other attempts [45, 46, 47, 48, 49] the BFKL equation has been modified in the infrared sector

in order to model some non-perturbative effects and at least try to avoid the strong inconsistency one has using the BFKL evolution which does not separate perturbative and non-perturbative scales. This fact is evident on considering the diffusion mechanism of the gluon distribution in the transverse momentum space during the evolution in the rapidity leading to a typical normal distribution in $\ln \mathbf{k}^2$. In all these phenomenological models a better behaviour also due to some decrease of the pomeron intercept to more reasonable values was found, but one of the fundamental properties shown by the QCD in the Regge kinematical limit, that the gluon reggeizes, is lost.

The idea to introduce a running coupling in a manner consistent with the gluon reggeization has been proposed by M.Braun [13, 14] and in this way one is able to guarantee that the production amplitude in the one reggeized gluon exchange approximation, which serves as an input in the BFKL theory, satisfy unitarity in the leading order [22]. Thereby the whole scheme becomes self-consistent: otherwise one should add to the input amplitudes corrections following from the unitarity. Since the bootstrap condition, as discussed in section 1.x, is a crucial element for the reggeization of the gluon and for the theory of the reggeized gluons as a whole the only way to introduce the running coupling constant in a manner compatible with the gluon reggeization is to preserve the bootstrap. A second basic ingredient is the requirement that the outgoing gluon distribution has the same asymptotic behavior as the one predicted by the DGLAP equations (describing the renormalization group flow) in the DLLA.

In section 1 we present the basic equations in the form suitable for numerical analysis [15] for the cases $q = 0$ (forward scattering) and $q \neq 0$. After that we describe the method of the solution and present the numerical results for the intercept, slope and the wave function at $q = 0$. The results for the intercept and slope, on the whole, agree with those found in [13, 14] by the variational approach. An interesting new result is the existence of a second pomeron with the intercept minus one, roughly speaking, two time less than for the leading one, but still positive. We end this section applying these results to the study of the asymptotical behaviour of the cross-section for the $\gamma^* \gamma^*$ scattering, including some unitarization effects due to multipomeron scattering by mean of an eikonal formulation valid in the large N_c limit [50], and try also to extrapolate to the hadron case.

Therefore the results of our study show that with the running coupling included the pomeron equation possesses bound state solutions which have negative energy and thus intercepts greater than unity. These solutions correspond to supercritical pomerons in the old sense, that is, they represent simple poles in the complex angular momentum plane. However the sub-dominant pomeron does not seem to play an important role in describing the asymptotical behaviour of the amplitudes. The intercept of the leading pomeron singularity depend weakly on the infrared regulator parameter and stay in the region 0.35–0.5 for its physically reasonable values. The introduction of the running coupling and thus a scale provides for a nontrivial slope for the pomeron, which is responsible for the physically reasonable behaviour of the cross-sections at very high energies.

For realistic photonic cross-sections and, with a rather crude approximation, for the hadronic ones the estimated unitarization effects begin to be felt at extraordinary high energies, of the order 100–1000 TeV (or equivalently $x < 10^{-10} - 10^{-12}$). Until these energies a single pomeron

exchange remains a very good approximation to the asymptotic amplitude. By the way we remember that no effects due to triple pomeron interaction are here taken into account and these are seen to be required as shown by the unitarization program (see the third chapter). Assuming that the picture is not completely changed by the last consideration, a comparison to the experimental cross-sections and structure functions at the highest energy (lowest x) achieved seems to confirm the widespread opinion that we are still rather far from the asymptotical regime and that other states, different from the supercritical pomerons, give the dominant contribution.

In section 2 we apply our model with the two found supercritical pomerons to jet production [16]. This process has been extensively studied in the framework of the standard BFKL approach with a fixed coupling [51, 45]. As is well-known, this analysis has lead to some far-reaching conclusions as to the importance of mini-jet production at high energies and the logarithmic rise of the multiplicity. However in the fixed coupling approach, various *ad hoc* modifications of the basic BFKL model had to be introduced to cut off the spectrum at low $k_\perp$ and also to correctly reproduce the high $k_\perp$ tail of the spectrum. Both these problems are naturally resolved by the introduction of a running coupling in our model, in which no new parameters appear in contrast to the fixed coupling approach. We shall study the asymptotic behaviour of the jet production cross section using the asymptotic form of the pomeron wave functions shown in appendix A.1 and present some numerical calculations.

Since we feel that to describe the present experimental data it is necessary to take into account all the states from the the spectrum of pomeron equation, this one has been converted into an evolution equation in $1/x$ and solved with an initial condition at some (presumably small) value $x = x_0$. In such an approach [17], taking a non-perturbative input at $x = x_0$ adjusted to the experimental data, also the problem of coupling the pomeron to the hadronic target is solved in an effective way. This approach is shown in details in section 3 where we state the basic equations and we try to pass from the gluon density to the observable structure function.

2.1 Pomeron equation with running coupling

Let us consider the bootstrap condition for the reggeization of the gluon:

$$[\omega(q) - \omega(q_1) - \omega(q_2)]\phi(q_1) = \int \frac{d^2 q'_1}{(2\pi)^2} K_q(q_1, q'_1) \phi(q'_1) \quad (2.1)$$

This is in general a very complicated functional integral equation which relates ω , K and ϕ . Looking for a particular class of solutions of this equation (we look for ω and K , limiting ourselves to constant ϕ) we can start to write them making use of more than one arbitrary function. In fact we can write the reggeized gluon trajectory

$$\omega(q) = -\frac{N_c}{2} \int \frac{d^2 q_1}{(2\pi)^2} \frac{f_A(q)}{f_B(q_1) f_C(q_2)}, \quad q = q_1 + q_2 \quad (2.2)$$

30 Hard pomeron with a running coupling constant: a phenomenological approach

and the gluon pair interaction kernel in the gluon channel

$$K_q(q_1, q'_1) = \frac{N_c}{2} \left(\frac{f_A(q_1)}{f_B(q'_1)} + \frac{f_A(q_2)}{f_B(q'_2)} \right) \frac{1}{f_C(q_1 - q'_1)} - \frac{f_A(q)}{f_B(q'_1)f_C(q'_2)} \quad (2.3)$$

One must note that there are some constraints on the kernel K which must be satisfied. The first one is that K should be symmetric in the two gluons (indices 1 and 2) so we must require that $f_B = f_C$ from the last term in the kernel. The second requirement is that K should be symmetric in the initial and final gluons (quantities with momentum not primed and primed), after separating the two "gluon propagators" ($\frac{1}{f_B(q'_1)f_B(q'_2)}$). From the first term in the kernel we must thus require this symmetry for the expression

$$\frac{f_A(q_1)f_B(q'_2) + f_A(q_2)f_B(q'_1)}{f_C(q_1 - q'_1)} \quad (2.4)$$

Matching the expression with the corresponding one where the primed quantities are exchanged with the not primed we finally get $f_A = f_B$. So we arrive at $f_A = f_B = f_C = \eta$. The standard BFKL theory is characterized by $\eta = q^2/(2\alpha_s)$ and in general the gluon trajectory and the interaction kernel are correlated in terms of the η function.

Let us write the equation for the amplitude in the vacuum channel in the form from which the typical cancellation of the infrared divergences in the pure BFKL theory is manifest

$$(j-1)\phi(q_1) = N_c \int \frac{d^2 q'_1}{(2\pi)^2} \left[\frac{\eta(q_1)}{\eta(q_1 - q'_1)} \left(\frac{\phi(q'_1)}{\eta(q'_1)} - \frac{\phi(q_1)}{\eta(q'_1) + \eta(q_1 - q'_1)} \right) + (1 \leftrightarrow 2) - \frac{\eta(q)}{\eta(q'_1)\eta(q'_2)} \phi(q'_1) \right] \quad (2.5)$$

The idea in [13, 14] was to change $\eta(q)$ so that it corresponds to a running rather than to a fixed coupling. For the running coupling some conclusions about the form of $\eta(q)$ can be made considering the vacuum channel equation in the limiting case of very large q . In this case one has, assuming that $\eta(q)$ grows with q , the approximated equation in the forward direction ($q = 0$)

$$(j-1)\phi(q_1) = 2N_c \int \frac{d^2 q'_1}{(2\pi)^2} \theta(q_1^2 - q'^2_1) \frac{\phi(q'_1)}{\eta(q'_1)} \quad (2.6)$$

and in the leading approximation one gets, after an inverse Mellin transformation, an evolution equation for the fully amputated wave function

$$\frac{\partial}{\partial \ln q_1^2} \frac{\partial}{\partial \ln 1/x} \phi(x, q_1) = \frac{N_c}{2\pi} \frac{q_1^2}{\eta(q_1)} \phi(x, q_1) \quad (2.7)$$

which, on comparing with the DGLAP evolution equation in the leading order in $\ln 1/x$ (that is, in the double leading log approximation (DLA)) for the gluon distribution,

$$\frac{\partial}{\partial \ln q_1^2} \frac{\partial}{\partial \ln 1/x} xg(x, q_1) = N_c \frac{\alpha_s(q_1^2)}{\pi} xg(x, q_1) \quad (2.8)$$

allows one to find the asymptotic form of $\eta(q)$:

$$\eta(q) \simeq \frac{q^2}{2\alpha_s(q^2)}, \quad q \rightarrow \infty \quad (2.9)$$

and a proportionality relation between the amputated pomeron wave function $\phi(x, k^2)$ and the gluon distribution $xg(x, k^2)$. It evidently differs from the fixed coupling case by changing the fixed coupling constant α_s to a running one $\alpha_s(q^2)$. As a result, with a running coupling, both the gluon trajectory and its interaction have to be changed simultaneously in an interrelated manner, so that the resulting equation is different from the BFKL one already in the leading order.

The behaviour of $\eta(q)$ at small q , comparable or even smaller than the QCD parameter Λ , cannot be established from any theoretical calculation, since this domain is non-perturbative. To take into account these confinement effect we choose $\eta(q)$ at finite q in a simple form:

$$\eta(q) = (b/2\pi)f(q) \quad ; \quad f(q) = (q^2 + m_1^2) \ln((q^2 + m^2)/\Lambda^2) \quad (2.10)$$

with $b = (1/4)(11 - (2/3)N_F)$ and $m \geq \Lambda$, which agrees with (2.9) for large q and remains finite up to $q = 0$. It allows for the freezing of the coupling and the confinement proper to occur at somewhat different scales (m and m_1 respectively). However, on physical grounds, one feels that they should be of the same order.

A preliminary study of the properties of the pomeron with $\eta(q)$ given by (2.10) for $m = m_1$ was performed by the variational technique in [13, 14]. It was found that the intercept depended on the ratio m/Λ quite weakly: as m/Λ changes from 1.5 to 5.0 the intercept (minus one) Δ falls from 0.4 to 0.25. On the other hand, the slope depends on this ratio very strongly. This allows to fix the ratio m/Λ to values in the interval $3.0 \div 4.0$.

This variational study, although very simple, cannot however give values for the intercept and especially for the slope with some precision. Still less can be found by this method about the properties of the pomeron wave function essential for the high-energy behaviour of the physical amplitudes, for some aspects studied in [50]. Finally, one does not receive any knowledge about the existence of other solutions with a positive intercept. All these reasons give us a motivation to undertake a numerical study of the two-gluon vacuum channel equation with in general the gluon trajectory and interaction given by

$$\begin{aligned} \omega(q) &= -\frac{N_c}{2} \int \frac{d^2 q_1}{(2\pi)^2} \frac{\eta(q)}{\eta(q_1)\eta(q_2)} \\ K_q(q_1, q'_1) &= -T_1 T_2 \left(\left(\frac{\eta(q_1)}{\eta(q'_1)} + \frac{\eta(q_2)}{\eta(q'_2)} \right) \frac{1}{\eta(q_1 - q'_1)} - \frac{\eta(q)}{\eta(q'_1)\eta(q'_2)} \right) \end{aligned} \quad (2.11)$$

where T is the gluon colour vector ($T_1 T_2 = -N_c$ in the vacuum channel) and the function $\eta(q)$ satisfying (2.9).

The parametrization (2.10), with b fixed, does not evidently take into account the change in N_F which occurs on flavour thresholds. In principle this can easily be remedied by assuming N_F and thus b depending on q^2 and varying accordingly. Our theory possesses a scale (actually

several scales, m, m_1 and Λ), so that the asymptotic behaviour (5) used to determine η refers to $q^2 \gg m^2, m_1^2, \Lambda^2$. Since $m \sim m_1$ is of the order of 1 GeV (the actual choice we make in following is $m = m_1 = 0.82 \text{ GeV}$), the only thresholds to be accounted for are those of the heavier quarks, b and t . However one should take into account that our theory is much better behaved than the scale-less BFKL model. In particular, in our case the pomeron possesses a normalizable wave function. So, in contrast to the BFKL case, the characteristic momenta which form the pomeron are determined by the same scale $m \sim m_1$ and are therefore of the order of several GeV/c . For this reason we expect to find the pomeron properties with sufficient precision fixing $N_F = 4$ and forgetting about b and t thresholds. Accordingly, the calculations presented are performed with a constant $N_F = 4$. Clearly the found momentum wave function is strictly speaking valid up to momenta squared of the order $100 (\text{GeV}/c)^2$, when the beauty threshold should be taken into account to slightly lower the value of our parameter b (by 8%). In applications to the structure functions this restricts the domain of applicability of the calculated pomeron wave function to $Q^2 \leq 100 (\text{GeV}/c)^2$.

2.1.1 Basic equations. Pomeron at $q = 0$

We consider the physical case $N_c = 3$. The units are chosen to have $\Lambda = 1$. In relating to observable quantities we take $\Lambda = 0.2 \text{ GeV}$.

The pomeron equation is the eigenvalue equation

$$\left(-\omega(q_1) - \omega(q_2)\right)\phi(q_1) + \int \frac{d^2 q'_1}{(2\pi)^2} K_q^{vac}(q_1, q'_1)\phi(q'_1) = E(q)\phi(q_1) \quad (2.12)$$

where the "energy" eigenvalue $E(q)$ is related to the pomeron trajectory via

$$\alpha(q) = 1 - E(q) \simeq 1 + \Delta - \alpha' q^2 \quad (2.13)$$

The last equation, well satisfied for small q , defines the intercept Δ and the slope α' . In (2.12) the trajectories ω and the kernel K^{vac} are given by the eqs. (2.11) with $T_1 T_2 = -3$ and the function η given by (2.10). To symmetrize the kernel we pass to the function

$$\psi(q_1) = \phi(q_1) / \sqrt{\eta(q_1)\eta(q_2)} \quad (2.14)$$

We also take out the common numerical factor $6/((11 - 2/3N_F)\pi)$ and express all terms via the function $f(q)$ also defined in (2.10). Then the equation for ψ takes the form

$$A_q(q_1)\psi(q_1) + \int d^2 q'_1 L_q(q_1, q'_1)\psi(q'_1) = \epsilon(q)\psi(q_1) \quad (2.15)$$

Here the "kinetic energy" is

$$A_q(q_1) = (1/2) \int \frac{d^2 q'_1 f(q_1)}{f(q'_1) f(q_1 - q'_1)} + (1/2) \int \frac{d^2 q'_2 f(q_2)}{f(q'_2) f(q_2 - q'_2)} \quad (2.16)$$

The interaction kernel consists of two parts, a quasi-local and a separable one: $L = L^{(ql)} + L^{(sep)}$

They are given by

$$L_q^{(ql)}(q_1, q'_1) = -\sqrt{\frac{f(q_1)}{f(q_2)}} \frac{1}{f(q_1 - q'_1)} \sqrt{\frac{f(q'_2)}{f(q'_1)}} - \sqrt{\frac{f(q_2)}{f(q_1)}} \frac{1}{f(q_2 - q'_2)} \sqrt{\frac{f(q'_1)}{f(q'_2)}} \quad (2.17)$$

and

$$L_q^{(sep)}(q_1, q'_1) = \frac{f(q)}{\sqrt{f(q_1)f(q_2)f(q'_1)f(q'_2)}} \quad (2.18)$$

Both parts are evidently symmetric in q_1 and q'_1 . The scaled energy ϵ is related to the initial one by

$$E = \frac{6}{\pi(11 - (2/3)N_F)} \epsilon \quad (2.19)$$

eq. (2.15) simplifies in the case when the total momentum of the two gluons is equal to zero. With $q = 0$ the two parts of the kinetic energies become equal and the square roots in (2.17) turn to unity. So at $q = 0$ the equation retains its form (2.15) with

$$A_0(q_1) = \int \frac{d^2 q'_1 f(q_1)}{f(q'_1) f(q_1 - q'_1)} \quad (2.20)$$

and the interaction where now

$$L_0^{(ql)}(q_1, q'_1) = -\frac{2}{f(q_1 - q'_1)} \quad (2.21)$$

has really become local and

$$L_0^{(sep)}(q_1, q'_1) = \frac{f(0)}{f(q_1)f(q'_1)} \quad (2.22)$$

This is the equation which we shall solve numerically.

To reduce to a one-dimensional problem we introduce the angular momentum of the gluons n and choose the solution in the form

$$\psi(q) = \psi_n(q^2) \exp in\phi \quad (2.23)$$

where ϕ is the azimuthal angle. Integrating over it in the equation, we obtain an one-dimensional integral equation for the radial function $\psi_n(q^2)$:

$$A_0(q)\psi_n(q^2) + \int dq_1^2 L_n(q^2, q_1^2)\psi(q_1^2) = \epsilon\psi_n(q^2) \quad (2.24)$$

with the kernel now given by

$$L_n(q^2, q_1^2) = -B_n(q^2, q_1^2) + \delta_{n0}\pi \frac{f(0)}{f(q)f(q_1)} \quad (2.25)$$

where

$$B_n(q^2, q_1^2) = \int_0^{2\pi} d\phi \frac{\cos n\phi}{f(q^2 + q_1^2 - 2qq_1 \cos \phi)} \quad (2.26)$$

Note that A_0 can be expressed via B_0 :

$$A_0(q) = (1/2) \int dq_1^2 B_0(q^2, q_1^2) \frac{f(q)}{f(q_1)} \quad (2.27)$$

Evidently eq. (2.24) is very similar to a Schrödinger equation with an attractive interaction provided by the local term and a positive kinetic energy described by A , which however grows very slowly at high momenta (as $\ln \ln q$). Therefore the attraction becomes smaller with growing n and we expect to find negative energies, corresponding to intercepts larger than unity, only for small n . Remember that for the BFKL pomeron only the isotropic state with $n = 0$ has a negative energy. Our calculations reveal that the introduction of the running coupling following does not change this situation: states with $|n| > 0$ all have positive energies. So in the following we consider the case $n = 0$.

2.1.2 Pomeron at $q \neq 0$: the slope

With $q \neq 0$ the pomeron equation becomes essentially two dimensional. Rather than attempt to solve it numerically at all q we limit ourselves to small values of q and determine not the whole trajectory $\alpha(q)$ but only the slope α' defined by (2.13). This can be done in a much simpler manner using a perturbative approach. We present "the Hamiltonian" in (2.15) $H_q = A_q + L_q$ in the form $H_q = H_0 + W(q)$ and calculate analytically $W(q)$ up to terms of the second order in q . Then for small q the value of the energy $\epsilon(q)$ will be given by the standard perturbation formula

$$\epsilon(q) = \epsilon(0) + \langle W(q) \rangle \quad (2.28)$$

where $\langle \rangle$ means taking the average with the wave function at $q = 0$, determined from the numerical solution of the equation discussed in the previous section. Thus we avoid solving the two-dimensional problem, but, of course, cannot determine more than the slope. Fortunately it is practically all we need to study the high-energy asymptotics (although, of course, the knowledge of the trajectory as a whole might be of some interest).

In order to derive an expression for $W(q)$ we pass to the relative momenta l and l' $q_{1(2)} = (1/2)q + (-)l$ and similarly for the primed momenta. Up to the second order in q we have

$$f(q_1) = f(l) [1 + a_1(q \cdot l) + \frac{a_1}{4} q^2 + \frac{a_2}{2} (q \cdot l)^2] \quad (2.29)$$

where

$$a_1 = a_1(l) = [1 + \ln(l^2 + m^2) - \frac{m^2 - m_1^2}{l^2 + m^2}] \frac{1}{f(l)} \quad (2.30)$$

$$a_2 = a_2(l) = [\frac{1}{l^2 + m^2} + \frac{m^2 - m_1^2}{(l^2 + m^2)^2}] \frac{1}{f(l)} \quad (2.31)$$

The expansion for $f(q_2)$ differs by changing the sign of l (or of q); for $f(q'_1)$ and $f(q'_2)$ it suffices to replace l by l' in the expressions for $f(q_1)$ and $f(q_2)$. We use the notation $a'_1 = a_1(l')$ and

$a'_2 = a_2(l')$. We also need the expansion for $f(q)$:

$$f(q) = f(0)(1 + a_3 q^2) + O(q^4) \quad ; \quad a_3 = \left(\frac{m_1^2}{m^2} + \ln m^2 \right) \frac{1}{f(0)} \quad (2.32)$$

The perturbation $W(q)$, up to second order in q , can be expressed via the introduced functions $a_{1,2}$ and $a'_{1,2}$ and the constant a_3 . After some calculations we find a part of W coming from the kinetic term in Hamiltonian in the form

$$W_1(l) = \frac{1}{2} \int d^2 l' \frac{f(l)}{f(l')} \left\{ \left(\frac{1}{f(l-l')} + \frac{1}{f(-l-l')} \right) \left[-a'_1(q \cdot l') - \frac{a'_1}{4} q^2 + (a_1'^2 - \frac{a'_2}{2})(q \cdot l')^2 \right] + \left[-\frac{a_1 a'_1(q \cdot l)(q \cdot l')}{f(l-l')} + \frac{a_1 a'_1(q \cdot l)(q \cdot l')}{f(-l-l')} \right] \right\} + A_0(l) \left[\frac{a_1}{4} q^2 + \frac{a_2}{2} (q \cdot l)^2 \right] \quad (2.33)$$

The part of W coming from the quasi-local part of the interaction can be written as

$$W_2(l, l') = \frac{1}{2} [a_1(q \cdot l) - a'_1(q \cdot l')]^2 L_0^{(ql)}(l, l') \quad (2.34)$$

and the one coming from the separable part as

$$W_3(l, l') = \left[(a_3 - \frac{a_1 + a'_1}{4}) q^2 - \frac{1}{2} (a_2 - a_1^2) (q \cdot l)^2 - \frac{1}{2} (a'_2 - a_1'^2) (q \cdot l')^2 \right] L_0^{(sep)}(l, l') \quad (2.35)$$

As mentioned, only isotropic solutions have the intercept larger than one and are of interest. Then the expression for $W(q) = \sum_{i=1,2,3} W_i$ has to be integrated over the azimuthal angles. Thus integrated values will be denoted $\hat{W}_i$, $i = 1, 2, 3$. Using (2.25)-(2.27), they can be conveniently expressed via the kernel B_n (eq. (2.26)):

$$\frac{\hat{W}_1}{2\pi q^2} = \frac{1}{4} \int dl'^2 \frac{f(l)}{f(l')} \left\{ \left[-\frac{a'_1}{2} + (a_1'^2 - \frac{a'_2}{2}) l'^2 \right] B_0(l, l') - a_1 a'_1 l l' B_1(l, l') \right\} + \frac{A_0(l)}{4} (a_1 + a_2 l^2) \quad (2.36)$$

$$\frac{\hat{W}_2}{2\pi q^2} = -\frac{1}{2} (a_1^2 l^2 + a_1'^2 l'^2) B_0(l, l') + a_1 a'_1 l l' B_1(l, l') \quad (2.37)$$

$$\frac{\hat{W}_3}{2\pi q^2} = \frac{f(0)}{f(l)f(l')} \left[2\pi (a_3 - \frac{a_1 + a'_1}{4}) - \frac{1}{4} l^2 (a_2 - a_1^2) - \frac{1}{4} l'^2 (a'_2 - a_1'^2) \right] \quad (2.38)$$

The slope is given by the momentum average of the sum of these expressions, taken with a given isotropic wave function:

$$\alpha' = -\frac{(1/2) \int d^2 l d^2 l' \psi(l) \psi(l') (\hat{W}_2 + \hat{W}_3) + \int d^2 l \psi(l)^2 \hat{W}_1}{2\pi q^2 \int d^2 l \psi^2(l)} \quad (2.39)$$

2.1.3 Numerical procedure and results

Eq. (2.24) was first changed to the variable $t = \ln q^2$ whereupon the wave function and the kernel transform according to

$$\psi(q^2) \rightarrow \tilde{\psi}(t) = q \psi(q^2) \quad (2.40)$$




Figure 2.1: .Pomeron intercepts as a function of the infrared regulator mass $m = m_1$; $\Lambda = 0.2 \text{ GeV}$ (a) and for different values of the confinement parameter m_1 and the coupling freezing parameter m (b).

and

$$L(q^2, q_1^2) \rightarrow \tilde{L}(t, t_1) = qq_1 L(q^2, q_1^2) \quad (2.41)$$

Performing a discretization on a grid the equation was reduced to a finite system of linear equations by approximating the integral by a sum

$$\int_{-\infty}^{\infty} dt F(t) \simeq \sum_{i=1}^n w_i F(t_i) \quad (2.42)$$

with the grid points t_i and weights w_i depending on the chosen approximation scheme. The final equation is thus

$$\sum_{j=1}^n B_{ij} x_j = \epsilon x_i, \quad i = 1, \dots, n \quad (2.43)$$

where

$$x_i = \sqrt{w_i} \tilde{\psi}(t_i) \quad (2.44)$$

and

$$B_{ij} = A(t_i) \delta_{ij} + \sqrt{w_i w_j} \tilde{L}(t_i, t_j) \quad (2.45)$$

The vector space for $\mathbf{x}$ has been chosen in order to obtain a symmetric matrix B . After determining the lowest eigenvalues ϵ and the corresponding eigenvectors x_i the wave function in the momentum space is directly given by (2.40) and (2.44) at points $q^2 = \exp t_i$. It should be normalized according to

$$\int \frac{d^2 q}{(2\pi)^2} |\psi(q)|^2 = 1 \quad (2.46)$$

Note that this wave function is a partially amputated one (see eq. (2.14)). The full (non-amputated) wave function is given by $\Phi(q) = \psi(q)/\eta(q)$ (in the forward case). It is this function that appears in the physical amplitudes.

The results for the lowest (and negative) eigenvalues of energy for the case $n = 0$ (isotropic




Figure 2.2: Pomeron slopes as a function of the infrared regulator mass $m = m_1$; $\Lambda = 0.2 \text{ GeV}$.

pomeron) are presented in Fig. 2.1. Actually energies with an opposite sign are shown, which according to (2.13) give precisely the intercepts (minus one). As mentioned, the QCD scale here and in the following is taken to be $\Lambda = 0.2 \text{ GeV}$. In Fig. 2.1.a the intercepts are shown for the case when the two scales m and m_1 in (2.10) are equal. Fig. 2.1.b illustrates the dependence of the intercepts on the ratio m/m_1 . The most interesting observation which follows from these figures at once is that in all cases one observes two positive intercepts, which correspond to two different supercritical pomerons, the leading and subleading ones. The single cut, characterizing the BFKL spectrum, is deformed with the introduction of the running coupling and a discrete part of the spectrum, giving some poles, appears. The intercept of the leading pomeron is found to be in accordance with the earlier calculations [13, 14], performed by the variational method (it is slightly larger, which was to be expected). For physically realistic values of m and m_1 in the interval $0.5 \div 1.0 \text{ GeV}$ it takes on values in the region of $0.5 \div 0.3$ falling with the masses m and m_1 . The same trend is seen for the intercepts of the subleading pomeron, which lie in the interval $0.25 \div 0.15$.

The slopes of the two found pomerons are shown in Fig. 2.2 as a function of m for the case $m = m_1$. The slopes depend very strongly on the value of the regulator mass. The physically reasonable slopes for the dominant pomeron of the order of $\alpha' \sim 0.25 \text{ (GeV/c)}^{-2}$ restrict the values of m to the region $0.7 - 0.9 \text{ GeV}$. So finally we choose

$$m = 0.82 \text{ GeV} \quad (2.47)$$

which leads to the following parameters of the leading (0) and subleading (1) pomerons

$$\Delta_0 = 0.384, \quad \alpha'_0 = 0.250 \text{ (GeV/c)}^{-2}; \quad \Delta_1 = 0.191, \quad \alpha'_1 = 0.124 \text{ (GeV/c)}^{-2} \quad (2.48)$$

In Fig. 2.3 we show the coordinate space wave functions $\Phi(r)$ of these two pomerons.




Figure 2.3: Coordinate space wave functions for the leading ($\Phi_0(r)$) and subleading ($\Phi_1(r)$) pomerons. Both r and Φ are in units $1/\Lambda \simeq 1 \text{ fm}$.

2.1.4 Pomeron and the high-energy scattering.

To apply the found results to the actual physical processes one has to couple the pomerons to the external sources corresponding to the colliding particles. The only way to do it in a more or less rigorous manner is to assume that both the projectile and target are highly virtual photons, or heavy onia, with momenta q and p respectively, $-q^2 = Q^2 \gg \Lambda^2$ and $-p^2 = P^2 \gg \Lambda^2$. Then the non-perturbative effects inside the target and projectile can safely be neglected. We shall address to the very complicated problem of coupling the gluons to the virtual photons in the running coupling framework in section 2.3 and consider here the simple lowest order approach, sufficient for the BFKL pomeron, where the coupling is small and fixed. This corresponds to taking the contribution of a pure quark loop into which the incoming photon goes. We shall then need of the colour densities of the virtual photons whose explicitly form, in the forward case, was found in [46] for both transverse and longitudinal photons. In the appendix A.2 we present a generalization for the non-forward case, both in momentum and coordinate representation, since we shall need of it, at least formally, in the third chapter, where we return to the fixed coupling formalism.

We shall try in the following to understand the role of the two found pomeron states and analyze the situation in which more than one pomeron is exchanged. The last situation can be seen as an attempt to understand some aspects which are related to higher order corrections needed to restore the unitarity. Incidentally the true situation is more complicated, as is shown in chapter 3, since, for a two pomeron amplitude, one has to take into account the triple pomeron interaction, which can take a form which absorbs completely the double pomeron exchange.

Nevertheless the contribution of multipomeron scattering to some processes is of interest and we shall try to analyze it as done in [50] where the scattering amplitude in the large colour number limit takes an eikonal form for fixed transverse dimensions of the projectile and target




Figure 2.4: Structure functions of a virtual photon ($P = 1 \text{ GeV}/c$) at $Q^2 = 4$ and $100 (\text{GeV}/c)^2$ as a function of x (a) and of the proton at $Q^2 = 4$ and $100 (\text{GeV}/c)^2$ as a function of x (b) (solid curves). Dashed curves show the contribution of a single pomeron exchange.

and leads to a cross-section

$$\sigma = 2 \int d^2 R d^2 r d^2 r' \rho_q(r) \rho_p(r') \left(1 - e^{-z(\nu, R, r, r')}\right) \quad (2.49)$$

where

$$z(\nu, R, r, r') = (1/8) \int \frac{d^2 q d^2 q_1 d^2 q'_1}{(2\pi)^6} G(\nu, q, q_1, q'_1) e^{iqR} \prod_{i=1,2} (1 - e^{iq_i r}) (1 - e^{iq'_i r'}) \quad (2.50)$$

is essentially the Fourier transform of the (non-amputated) Green function of eq. (2.12), $G(\nu, q, q_1, q'_1)$, considered as a function of the energetic variable $\nu = pq$ and with $q = q_1 + q_2 = q'_1 + q'_2$. The functions ρ_q and ρ_p correspond to the colour densities of the projectile and target photons, respectively.

The found supercritical pomerons represent a part of the total pomeron spectrum which contributes to the Green function, in the high energy limit, with a term given by

$$G_P(\nu, q, q_1, q'_1) = \sum_{i=0,1} \nu^{\alpha_i(q)-1} \Phi_i(q_1, q_2) \Phi_i^*(q'_1, q'_2) \quad (2.51)$$

where α_i and Φ_i are the trajectories and wave functions of the leading (0) and subleading (1) pomerons. At high ν we can neglect the dependence on the total momentum q of the wave functions, taking them at $q = 0$, and approximate the trajectories according to (2.13). Then all the quantities in (2.49) become determined, so that we can calculate the cross-sections for both the transversal and longitudinal projectile photon and thus find the structure function of the virtual photon target. We have taken for the latter a transversal photon with the lowest momentum admissible of $P = 1 \text{ GeV}/c$. The resulting structure functions are presented in Fig. 2.4.a for the interval of small x which we extended to extraordinary small values to clearly see the unitarization effects.

To move closer to reality one has to consider hadronic target and projectiles. The confinement effects make any rigorous treatment of such a case impossible. Rather than to introduce arbitrary parameters (in fact, functions) we extend the formula (2.49) to hadronic target and projectile substituting the photonic colour densities by hadronic ones. For the latter we take a Gaussian form and a normalization which follows from the comparison to the electromagnetic densities with only the simplest quark diagrams taken into account. In particular for the proton we take the Gaussian ρ , with the observed electromagnetic proton radius and normalized to three. Such a treatment, in all probability, somewhat underestimates the density, since it does not include coupling to gluons.

The proton structure functions and the proton-proton total cross-sections which follow from this approximation for the densities are shown in Figs. 2.4.b and 2.5 respectively. To see the unitarization effects we had again to consider extraordinary high values of $1/x$ and energies, well beyond our present experimental possibilities.

In discussing these results, we have first to note that their overall normalization is somewhat undetermined, since the exact scale at which ν enters into $\ln \nu$ factors is unknown. A second point to note is that the subleading pomeron contribution is always very small: it amounts to a few percent at smallest values of $1/x$ and s considered and naturally gets still smaller at higher $1/x$ or s .

As one observes from Figs. 2.4-2.5, the structure functions and cross-sections monotonously rise with $1/x$, s and Q^2 . Studying the asymptotics of the solutions of eq. (2.12) at high q and of eq. (2.49) one can show that this rise is logarithmic. In particular, the structure function of the virtual photon rises as $\ln^4(1/x)$ and as $\ln^\beta(q^2)$ with $\beta \sim 2.5$. The proton-proton cross-section eventually rise as $\ln^2 s$, as expected. Comparison to the Froissart bound (dash-dotted line in Fig. 7) shows however that it remains far from being saturated.

The most interesting result that follows from Figs. 2.4-2.5 is that the unitarization effects become visible only at exceedingly very small values of x or very large values of s , well outside the range of the present experiment. They appear earlier at lower Q^2 . Still at the smallest value $Q = 2 \text{ GeV}/c$ considered, the exchange of more than one pomeron achieves only 15% of the total for the proton structure function at $x = 10^{-10}$. Likewise the relative contribution of many pomerons to the proton-proton cross-section rises to 23% only at $\sqrt{s} \sim 10^5 \text{ GeV}$.

Comparing the calculated proton structure functions and the cross-sections with the experimental results at highest $1/x$ and s available we observe that our results are essentially smaller than the observed ones. Experimental value of $F_{2p}(Q^2, x)$ at $Q^2 = 8.5 (\text{GeV}/c)^2$ and $x = 0.000178$ is $1.19 \pm 0.05 \pm 0.16$ [52]. Our calculations only give a value 0.17. The $\bar{p}p$ cross-section at $\sqrt{s} = 1800 \text{ GeV}$ is around 80 mbn [53], whereas our result is 18.5 mbn . Of course, having in mind the uncertainties in the overall normalization and a very crude picture for the pomeron coupling to the proton assumed, one cannot ascribe too much importance to this fact. However one is tempted to explain this underestimation of the experimental values by the simple reason that we are too far from the pure asymptotic regime yet and that other solutions of eq. (2.12) different from the found supercritical pomerons and having their intercepts around unity give the bulk of the contribution at present energies. This may also explain the notorious discrepancy between a high value of the hard pomeron intercept, of the order 0.35–0.5, and the




Figure 2.5: Proton-proton total cross-sections as a function of c.m. energy $\sqrt{s}$ (the solid curve). The dashed curve shows the contribution of a single pomeron exchange. The dash-dotted curve marks the Froissart bound.

observed slow growth of the experimental cross-section, well described by the "soft pomeron" with an intercept around 0.08.

Forgetting for the moment the triple pomeron interactions, if this picture is correct then we may expect that with the growth of energy the cross-sections will grow faster and faster, until at $\sqrt{s} \sim 10 \text{ TeV}$ they will become well described by a pure hard pomeron with the intercept 0.35–0.5. This power growth will continue until energies of an order 1000 TeV when finally the unitarity corrections set in to moderate the growth in accordance with the Froissart bound.

2.2 Inclusive jet production.

2.2.1 General formalism.

We shall consider in the following the scattering of two highly virtual photons (or heavy "onia") to make the derivation more rigorous. To obtain the formula for the inclusive jet production let us recall the total cross section for one pomeron exchange [50]:

$$\sigma = \frac{1}{4} \int d^2r d^2r' \rho_q(r) \rho_p(r') \int \frac{d^2q_1 d^2q'_1}{(2\pi)^4} G(\nu, 0, q_1, q'_1) \prod_{i=1,2} (1 - e^{iq_i r})(1 - e^{iq'_i r'}) \quad (2.52)$$

Here $q(p)$ is the momentum of the projectile (target); $\nu = qp = (1/2)s$; $G(\nu, q, q_1, q'_1)$ is the non-amputated pomeron Green function in the momentum space, q being the total momentum of the two gluons and q_1 (q'_1) being the initial (final) momentum of the first gluon; $\rho_{q(p)}$ is the dipole colour density of the projectile (target). For the virtual photons we take the colour dipole densities from [44].

In the high energy limit the dominant term of the Green function comes from the two mentioned supercritical pomeron states (2.51) where α_i and Φ_i are the trajectories and wave functions of the leading (0) and subleading (1) pomerons.

Mini-jets appear as intermediate gluon states in the Green function (2.51). They possess




Figure 2.6: The substitution for the Green function used to calculate the inclusive jet production cross section.

arbitrary $k_\perp$ subject to condition $k_\perp^2 \ll s$. The inclusive cross-section for their production can be calculated by splitting the Green function in (2.52) as indicated in Fig. 2.6 whihc means that we perform the substitution

$$G(\nu, 0, q_1, q'_1) \Rightarrow \int \frac{d^2 k_1 d^2 k'_1}{(2\pi)^4} G(\nu_1, 0, q_1, k_1) V(k_1, k'_1) G(\nu_2, 0, k'_1, q'_1) \delta^{(2)}(k_1 + k_\perp - k'_1) \quad (2.53)$$

where, for the running coupling case,

$$V(k_1, k'_1) = 6 \frac{\eta(k_1) \eta(k'_1)}{\eta(k_1 - k'_1)} - 3\eta(0) \quad (2.54)$$

Remembering that $q_1 + q_2 = q'_1 + q'_2 = q = 0$ we obtain

$$I(y, k_\perp) \equiv \frac{d^3 \sigma}{dy d^2 k_\perp} = \int d^2 r d^2 r' \rho_q(r) \rho_p(r') \int \frac{d^2 q_1 d^2 q'_1}{(2\pi)^4} (1 - e^{iq_1 r}) (1 - e^{iq'_1 r'}) \int \frac{d^2 k_1 d^2 k'_1}{(2\pi)^4} \delta^{(2)}(k_1 + k_\perp - k'_1) G(\nu_1, 0, q_1, k_1) G(\nu_2, 0, k'_1, q'_1) [6 \frac{\eta(k_1) \eta(k'_1)}{\eta(k_1 - k'_1)} - 3\eta(0)] \quad (2.55)$$

The appropriate kinematical variables are the rapidity and transverse momentum of the observed gluon (jet), $y = \frac{1}{2} \ln \frac{k_+}{k_-}$ and $k_\perp$, respectively ; we also have $s_i = 2\nu_i$ and $s_{1(2)} =$

$k_\perp \sqrt{s} e^{-(+)y}$. It is convenient to use a mixed (momentum-coordinate) representation for the Green functions:

$$G(\nu, 0, r, k) = \int \frac{d^2 q_1}{(2\pi)^2} e^{i q_1 r} G(\nu, 0, q_1, k) \quad (2.56)$$

Defining $\Delta\Phi_i(r) = \Phi_i(r) - \Phi_i(0)$ we can write

$$G(\nu, 0, r, k) - G(\nu, 0, 0, k) = \sum_{i=0,1} \nu^{\Delta_i} \Delta\Phi_i(r) \Phi_i^*(k) \quad (2.57)$$

We also introduce

$$R_i^{q(p)} \equiv \langle \Delta\Phi_i \rangle_{q(p)} = \int d^2 r \rho_{q(p)}(r) \Delta\Phi_i(r) \quad (2.58)$$

In this notation and also using both the semi-amputated and full wave functions we find

$$I(y, k_\perp) = \sum_{i,j=0,1} R_i^q R_j^p \nu_1^{\Delta_i} \nu_2^{\Delta_j} \int \frac{d^2 k_1}{(2\pi)^4} \left[6 \frac{\psi_i(k_1) \psi_j(k_1 + k_\perp)}{\eta(k_\perp)} - 3\eta(0) \Phi_i(k_1) \Phi_j(k_1 + k_\perp) \right] \quad (2.59)$$

Since the found supercritical pomerons are isotropic in the transverse space the inclusive cross-section $I(y, k_\perp)$ also turns out to be isotropic and therefore we can integrate over azimuthal angles in (2.59). Defining for the full and semi-amputated wave functions the integrated quantities

$$\begin{aligned} \hat{\Phi}(k_1, k_\perp) &= \int_0^{2\pi} d\alpha \Phi(k_1 + k_\perp) \\ \hat{\psi}(k_1, k_\perp) &= \int_0^{2\pi} d\alpha \eta(k_1 + k_\perp) \Phi(k_1 + k_\perp) \end{aligned} \quad (2.60)$$

where α is the angle between k_1 and $k_\perp$, we finally obtain the inclusive cross section

$$\begin{aligned} \frac{d^2 \sigma}{dy dk_\perp^2} &= \frac{3}{4} \sum_{i,j=0}^1 e^{-y(\Delta_i - \Delta_j)} \left(\frac{k_\perp^2 s}{4} \right)^{\frac{1}{2}(\Delta_i + \Delta_j)} R_i^q R_j^p \\ &\quad \int \frac{dk_1^2}{(2\pi)^3} \left[2 \frac{\psi_i(k_1) \hat{\psi}_j(k_1, k_\perp)}{\eta(k_\perp)} - \eta(0) \Phi_i(k_1) \hat{\Phi}_j(k_1, k_\perp) \right] \end{aligned} \quad (2.61)$$

The results of numerical calculations of the cross-section (2.61) and also its generalization to more interesting cases of hadronic targets or/and projectiles will be discussed in the next section. In the rest of this section we shall study the asymptotic behaviour of the found inclusive jet production cross-section at very small and very large transverse momenta and also its y -dependence.

As to the latter, all y -dependence in (2.61) comes from the factor $\exp(-y(\Delta_i - \Delta_j))$ which has its origin in the existence of two different supercritical pomerons. Evidently in the limit $s \rightarrow \infty$ this dependence dies out, since the relative contribution of the sub-dominant pomeron becomes negligible. The model thus predicts an asymptotically flat y plateau at very high energies.

At small $k_\perp$ the cross-section (2.61) evidently goes down as $k_\perp^{\Delta_1}$, since all other factors are finite in this limit. However one should remember that (2.51) gives the dominant contribution

only while $(k_\perp^2 s)$ continues to be large. At too small $k_\perp$, when the above quantity becomes finite, all other states from the spectrum of the two-gluon equation (2.12), hitherto neglected, begin to give comparable or even dominant contribution, so that the found $k_\perp^{\Delta_1}$ behaviour ceases to be valid.

To find the asymptotic behaviour of the inclusive cross section for $k_\perp \rightarrow \infty$ we need to know the behaviour of the pomeron wave functions in the momentum space at $q \rightarrow \infty$ and in the ordinary space at $r \rightarrow 0$. In the appendix A.1 we show that

$$\begin{aligned} \psi(q) &\underset{q \rightarrow \infty}{\sim} \frac{1}{q^2} (\ln q^2)^\beta \\ \psi(r) &\underset{r \rightarrow 0}{\sim} \left(\ln \frac{1}{r}\right)^{\beta+1} \end{aligned} \quad (2.62)$$

where $\beta = -1 - \frac{3}{bE}$ so that in the forward scattering case β depends just on the intercept of the corresponding pomeron state. Let us study now the behaviour of (2.59) for $k_\perp \rightarrow \infty$. Using $\Phi(q) = \psi(q)/\eta(q) \sim \frac{1}{q^4} (\ln q^2)^{\beta-1}$ we find $\int d^2 q \Phi(q) < \infty$, so that for the second term in the integrand of (2.59) we get

$$\int d^2 k_1 \Phi_i(k_1) \Phi_j(k_1 + k_\perp) \xrightarrow{k_\perp^2 \rightarrow \infty} \Phi_j(k_\perp) \int d^2 k_1 \Phi_i(k_1) \sim \frac{1}{k_\perp^4} (\ln k_\perp^2)^{\beta_j-1} \quad (2.63)$$

To analyze the first term we use the identity

$$\int \frac{d^2 k_1}{(2\pi)^2} \psi_i(k_1) \psi_j(k_1 + k_\perp) = 2\pi \int_0^\infty r dr J_0(k_\perp r) \psi_i(r) \psi_j(r) \quad (2.64)$$

and the relation

$$\int_0^\infty r dr J_0(k_\perp r) f(r) = -\frac{1}{k_\perp} \int_0^\infty r dr J_1(k_\perp r) f'(r) \quad (2.65)$$

which is valid for naturally behaved f , such that $\left[r f(r) J_1(k_\perp r) \right]_{r=0}^{r=\infty} = 0$.

Putting $f(r) \sim (\ln \frac{1}{r})^{\beta_i + \beta_j + 2}$ we obtain the leading behaviour

$$\int d^2 k_1 \psi_i(k_1) \psi_j(k_1 + k_\perp) \sim \frac{1}{k_\perp^2} \left(\ln k_\perp^2 \right)^{\beta_i + \beta_j + 1} \quad (2.66)$$

So at $k_\perp \rightarrow \infty$ we find for the inclusive cross-section (2.59)

$$I(y, k_\perp) \sim \sum_{i,j=0,1} R_i^q R_j^p \left(\frac{k_\perp^2 s}{4} \right)^{\frac{\Delta_i + \Delta_j}{2}} e^{-y(\Delta_i - \Delta_j)} \frac{1}{k_\perp^4} \left(\ln k_\perp^2 \right)^{\beta_i + \beta_j} \quad (2.67)$$

This asymptotics corresponds to the standard quark-counting rules behaviour ($\sim 1/k_\perp^4$), modified by a power factor due the pomeron energetic dependence and a logarithmic factor coming from the pomeron wave function, i.e. from the running of the coupling. Note that in (2.67) the $k_\perp$ and s dependencies are separated. As a result we find that the average $\langle k_\perp \rangle$ is finite and independent of s and $\langle k_\perp^2 \rangle \sim s^\Delta$, since it formally diverges for (2.67) and one has to restrict $k_\perp^2 < s$.

For the the multiplicity

$$\langle n \rangle = \frac{1}{\sigma} \int \frac{d^3\sigma}{dyd^2k_\perp} dyd^2k_\perp \quad (2.68)$$

the standard asymptotic behaviour $\langle n \rangle = a \ln s + b$ is obtained. Indeed integrating (2.59) we get $\int dyd^2k_\perp \frac{d^3\sigma}{dyd^2k_\perp} = Bs^{\Delta_0} \ln s + Cs^{\Delta_0}$. Since the total cross-section has the form $\sigma = As^{\Delta_0}$ at large s , we get the mentioned asymptotical expression for the multiplicity. Of course this result is valid only in the extreme limit $s \rightarrow \infty$. At large but finite s the existence of two different pomerons leads to some additional non-trivial s -dependence.

It is instructive to compare our asymptotic results with those obtained in the BFKL fixed coupling model. In the latter case the inclusive cross section is of course badly behaved in the $k_\perp \rightarrow 0$ limit due to scale invariance. It is also very different in the high $k_\perp$ limit. At very large $k_\perp$ such that $\ln k_\perp \sim \sqrt{\ln s}$ one finds

$$\left(\frac{d^2\sigma}{dydk_\perp^2} \right)_{BFKL} \underset{k_\perp \rightarrow \infty}{\sim} a(y) \frac{(k_\perp^2 s)^\Delta}{k_\perp^2} \frac{e^{-\ln^2 k_\perp^2 / a^2(y)}}{\ln k_\perp^2 \sqrt{\ln s}} \quad (2.69)$$

where $a^2 \sim (\ln s - 4y^2 / \ln s)$. Thus for large $k_\perp$ the BFKL cross section goes down faster than any power. Also one obtains that $\langle \ln k_\perp \rangle \sim \sqrt{\ln s}$ so that both $k_\perp$ and $k_\perp^2$ grow with s .

To bring these predictions in better correspondence with the physical reality, as mentioned, various modifications of this orthodox BFKL approach have been introduced. In particular in [51, 45] the fusion of gluons via "fan" diagrams was assumed to take place at high gluonic densities, which was considered as a way to partially restore the s-channel unitarity. Then, under some additional assumptions, a $1/k_\perp^4$ asymptotic behaviour similar to (2.67) was found. In our model such a behaviour naturally follows without introducing additional assumptions or imposing the unitarity restrictions.

2.2.2 Numerical results.

Taking the wave function evaluated numerically in [15] we have computed the cross section (2.61).

In Fig. 2.7.a we present $d^2\sigma/dydk_\perp$ for the process $\gamma^*\gamma^*$ (in units $c = 1$). We have chosen the projectile photon to have virtuality $Q = 5GeV/c$ and the target one to have virtuality $P = 1GeV/c$. The center of mass energy is $\sqrt{s} = 540GeV$.

Of course, processes involving hadronic targets or/and projectiles are much more interesting from the practical point of view. However these require some non-perturbative input for the colour densities of the colliding hadrons. A possible way to introduce it is evidently to convert eq. (2.12) into an evolution equation in $1/x$ and take initial conditions for it from the existing experimental data. Postponing this complicated procedure for future studies, we use here a simpler approach, taking for the hadron (proton) a Gaussian colour density with a radius corresponding to the observed electromagnetic one. Such an approximation, in all probability, somewhat underestimates the coupling of the hadron to the pomeron, since the coupling of the latter to constituent gluons is neglected. Nevertheless, we hope that it gives a reasonable estimate for the inclusive cross-section. Using this approach we get the inclusive jet production




Figure 2.7: Inclusive jet production cross sections for the processes $\gamma^*\gamma^*$ at $\sqrt{s} = 540\text{GeV}$ with $Q^2 = 25(\text{GeV}/c)^2$ and $P^2 = 1(\text{GeV}/c)^2$ (a), for γ^*p at $\sqrt{s} = 540\text{GeV}$ with $Q^2 = 25(\text{GeV}/c)^2$ (b) and for pp at the same c.m.s. energy (c).

cross-sections for the γ^*p and pp scattering shown in Fig. 2.7.b and 2.7.c respectively.

A common feature of jet production in all processes is that the cross-section reaches a maximum at $k_{\perp} \approx 1\text{ GeV}/c$ from which it monotonously goes down both for smaller and larger $k_{\perp}$.

In Fig. 2.8 the cross-section $d\sigma/dy$ integrated over $k_{\perp}$ in the interval $0.5 \div 20\text{ GeV}/c$ is presented for the process pp . The limitations in the numerical calculations of the wave functions do not allow us to study higher values of $k_{\perp}$, so that we cannot numerically reach the region where the asymptotical behaviour (2.67) is strictly valid.

In Fig. 2.9 we show jet multiplicities as a function of s for the three studied processes $\gamma^*\gamma^*$, γ^*p and pp . As one observes, their magnitude and behaviour prove to be quite similar.

We also tried to estimate the average $\langle k_{\perp} \rangle$. Unfortunately, although it exists according to (2.67), its value results very sensitive to the high momentum tail of the pomeron wave function, poorly determined from our numerical calculations. To avoid this difficulty we chose to calculate the average $\langle \ln k_{\perp}/\Lambda \rangle$ instead. This average at fixed y depends on s and y only due to the existence of two different pomerons, so that this dependence should die out at large enough s . In fact the found average $\langle \ln k_{\perp}/\Lambda \rangle$ turns out to be practically independent of y and very




Figure 2.8: Cross-sections $d\sigma/dy$ for the pp process at $\sqrt{s} = 540 \text{ GeV}$.




Figure 2.9: Multiplicities $\langle n \rangle$ as a function of the center of mass energy $\sqrt{s}$ for the processes $\gamma^*\gamma^*$ (the solid curve), γ^*p (the dashed curve) and pp (the dotted curve).

weakly dependent on s in the whole studied range of s and y , rising from 3.97 at $\sqrt{s} = 20 \text{ GeV}$ to 4.19 at $\sqrt{s} = 20 \text{ TeV}$. These values imply a rather high average $k_\perp$ rising from $10.6 \text{ GeV}/c$ at $\sqrt{s} = 20 \text{ GeV}$ to $13.2 \text{ GeV}/c$ at $\sqrt{s} = 20 \text{ TeV}$.

Our calculations show that the introduction of a running coupling constant on the basis of the bootstrap condition cures all the diseases of the orthodox BFKL approach for jet production. At high $k_\perp$ the cross-section becomes well-behaved and more or less in accordance with the expectations based on the quark counting rules. At small $k_\perp$ no singularity occurs, although contributions from other states is expected to dominate.

As to s - and y -dependence, our predictions are even simpler than in the BFKL approach, since the running of the coupling converts the branch point in the complex angular momentum plane, corresponding to the BFKL pomeron, into poles, of which only two are located to the right of unity and contribute at high energies. As a result at super-high energies, when only the dominant pomeron survives, the y -dependence completely disappears and the s -dependence reduces to the standard s^{Δ_0} factor. At smaller s some y - and extra s -dependence appears due

to the existence of two supercritical pomerons.

With all these refinements, some basic predictions of the BFKL theory are reproduced. The cross-section for mini-jet production rises fast and saturates the total cross-section as $s \rightarrow \infty$. Jet multiplicities rise logarithmically.

However $\langle k_{\perp} \rangle$ turns out to depend on s weakly and its calculated value results pretty high, of the order of 10-12 GeV/c . This should be contrasted to the experimentally observed much lower values of $\langle k_{\perp} \rangle$ rising with energy. A natural explanation of this discrepancy follows from the results [15] shown in the first section of this chapter in particular from the study of the structure functions and total cross-sections in our model: at present energies the contribution from the two supercritical pomerons only covers a part of the observed phenomena because we are still rather far from the real asymptotics. The bulk of the contribution comes from other states, which produce a much softer spectrum of particles. The observed rise of the $\langle k_{\perp} \rangle$ is then related to the dying out of these sub-asymptotical states. Our prediction is then that the rise of $\langle k_{\perp} \rangle$ should saturate at the level of 10-12 GeV/c .

This circumstance has also to be taken into account when discussing the numerical results presented in Figs. 6-9. Probably they also illustrate predictions for considerably higher energies than the present ones, at which, according to the estimate made in [15], they should account for $\sim 20\%$ of the observed spectra.

To obtain predictions better suited for present energies one should evidently take into account all states present in the spectrum of the pomeron equation (2.12) and not only the two supercritical pomerons. To realize this program an evolution equation in ν following from (2.12) seems to be an appropriate tool. As mentioned, it could also effectively take into account the non-perturbative effects related to the pomeron coupling to physical hadrons.

2.3 Evolution of the gluon density.

For the forward scattering amplitude the pomeron equation (2.12) for the isotropic wave function with an inhomogeneous term reads

$$(H - E)\psi = \psi_0 \quad (2.70)$$

Taking the Mellin transformation of (2.70) one converts it into an evolution equation in $1/x$:

$$\frac{\partial}{\partial \ln 1/x} \psi(x, k) = -H\psi(x, k) \quad (2.71)$$

which should be supplemented with an initial condition at some $x = x_0$

$$\psi(x_0, k) = \psi_0(k) \quad (2.72)$$

containing the non-perturbative input about the coupling to the hadronic target.

The physical interpretation of the pomeron wave function is provided by the fact that in the DLLA scheme eq. (2.71) reduces to an equation for the fully amputated function $\phi(x, k) =$

$\eta(k)\psi(x, k)$:

$$\frac{\partial}{\partial \ln k^2} \frac{\partial}{\partial \ln 1/x} \phi(x, k) = \frac{3\alpha_s(k^2)}{\pi} \phi(x, k) \quad (2.73)$$

which coincides with the standard equation for the unintegrated gluon density $xg(x, k^2)$ in the DLLA limit. In fact, this circumstance lies at the root of our method to introduce a running coupling into the scheme. Thus we may identify

$$\phi(x, k) = cxg(x, k) \quad (2.74)$$

The normalizing factor c cannot be determined from the asymptotic equation (2.73). We shall be able to fix it by studying the coupling of the pomeron to the incoming virtual photon in the next section.

2.3.1 Coupling to the virtual photon

Once the function ϕ proportional to the gluon density is determined, one has to couple it to the projectile particle to calculate observable quantities. In particular, to find the structure function of the target one has to couple the gluons to the incoming virtual photon, that is, to find the colour density $\rho(q, k)$ which connects the photon of momentum q to the gluon of momentum k . This problem is trivial within the BFKL approach with a fixed small coupling. Then it is sufficient to take the colour density in the lowest order $\rho_0(q, k)$, which corresponds to taking for it the contribution of a pure quark loop into which the incoming photon goes.

The problem complicates enormously when one tries to introduce a running coupling into ρ . Then one has to take into account all additional gluon and $q\bar{q}$ pair emissions which supply powers of the logarithms of transverse momenta. Apart from making the coupling run, they will evidently change the form of $\rho(q, k)$. Unfortunately the bootstrap relation can tell us nothing about the ultimate form of the colour density with a running coupling, which essentially belongs to the t -channel with a vacuum colour quantum number. So we have to find a different way to introduce a running coupling into ρ .

A possible systematic way to do this consists in applying to the photon-gluon coupling the DGLAP evolution equation. One may separate the colour density from the rest of the amplitude by restricting its rapidity range to some maximal rapidity $y_0 \sim \log Q^2$ (which, of course should be much smaller than the overall rapidity $Y \sim \log Q^2/x$). Then the kinematical region of $\rho(q, k)$ will admit the standard DGLAP evolution in Q^2 . Solving this equation one will find the quark density at scale Q^2 of the gluon with momentum k (i.e. essentially the structure function of the gluon with the virtuality k^2). This is exactly the quantity needed to transform the calculated gluon density created by the target into the observable structure function of the target. As a starting point for the evolution one may take the perturbative colour density ρ_0 at some low Q^2 when the logs of the transverse momenta might be thought to be unimportant.

This ambitious program, combining both evolution in $1/x$ and Q^2 , does not, however, look very simple to realize. As a first step, to clearly see the effects of the introduction of a running coupling preserving the gluon reggeization, we adopt a more phenomenological approach here, trying to guess a possible correct form for $\rho(q, k)$ on the basis of simple physical reasoning and




Figure 2.10: Forward amplitude for the scattering of a photon with a reggeized gluon..

also using the DLLA to fix its final form.

With a pure perturbative photon colour density one would obtain for the γ^*p cross-section

$$\sigma(x, Q^2) = \int \frac{d^2k \rho_0(q, k) \phi(x, k)}{(2\pi)^2 \eta^2(k)} \quad (2.75)$$

In fact, the projectile particle should be coupled to the full pomeron wave function ϕ/η^2 . From the physical point of view this expression is fully satisfactory for physical particles. However it is not for a highly virtual projectile.

To see this, we first note that for the forward amplitude our method of introducing a running coupling reduces to a very simple rule: the scale at which the coupling should be taken is given by the momentum of the emitted real gluon ($(k - k')^2$ in the upper rung in Fig. 2.10). Now take Q^2 very large and apply the DLLA approximation. Then the momenta in the ladder become ordered from top to bottom

$$Q^2 \gg k^2 \gg k'^2 \gg \dots$$

In this configuration, as can be traced from (2.70) and (2.75), all α_s 's acquire the right scale (i.e. corresponding to the DGLAP equation) except for the upper rung: $\alpha_s(k^2)$ appears twice. In order to clearly see why this happens let us explicitly write eq. (2.70) which describes the interactions in the ladder of Fig. 2.10 (the kinetic part given by the trajectories of the two reggeized gluons is denoted by T)

$$(E - T(k))\phi(k) = \phi_0(k) + 2 \int \frac{d^2k'}{(2\pi)^2} \frac{\eta(k)}{\eta(k')\eta(k - k')} \phi(k') \quad (2.76)$$

Constructing by iteration the solution, for example at the second order, one has

$$\phi^{(2)}(k) = \frac{2}{E - T(k)} \int \frac{d^2 k'}{(2\pi)^2} \frac{\eta(k)}{\eta(k - k')} \frac{2}{E - T(k')} \int \frac{d^2 k''}{(2\pi)^2} \frac{1}{\eta(k'')\eta(k' - k'')} \frac{\phi_0(k'')}{E - T(k'')} \quad (2.77)$$

and by taking the limit of the DLLA approximations one obtains

$$\phi^{(2)}(k) = \frac{2}{E - T(k)} \int \frac{d^2 k'}{(2\pi)^2} \frac{1}{\eta(k')} \frac{2}{E - T(k')} \int \frac{d^2 k''}{(2\pi)^2} \frac{1}{\eta(k'')} \frac{\phi_0(k'')}{E - T(k'')} \quad (2.78)$$

It is evident that after taking this limit there is no cancellation of one $\eta(k)$ term in (2.75). Therefore this defect can be understood if one notices that the upper gluon is, in fact, coupled to a virtual particle. If this particle were a gluon, then the interaction K_q in (2.11) would cancel one of the two $\alpha(k^2)$'s and substitute it by an α taken at the scale corresponding to its own virtuality. We assume that something similar should take place also for virtual quarks to which the gluon chain may couple. The scale of the particle momenta squared which enter the upper blob in Fig. 10 should have the order Q^2 (this is the only scale that remains after these momenta are integrated out). As a result the lowest order density should be rescaled according to

$$\rho_0(q, k) \rightarrow \frac{\alpha(Q_1^2)}{\alpha(k^2)} \rho_0(q, k) \quad (2.79)$$

where Q_1^2 has the same order as Q^2 .

The approximation we assume is that the substitution (2.79) is sufficient to correctly represent the photon colour density with a running coupling. We shall check its validity by studying the quark density which results from (2.79) in the DLLA approximation and comparing it with the known result based on the DGLAP equation.

Explicitly the zeroth order density ρ_0 has the following forms for the transverse (T) and longitudinal (L) photons (see e.g. [44] and do the integration in the quark loop momenta)

$$\rho_0^{(T)}(q, k) = \frac{3e^2}{8\pi^2} \sum_f Z_f^2 \int_0^1 d\alpha \left((\alpha^2 + (1 - \alpha)^2)((1 + 2z^2)g(z) - 1) + \frac{\zeta}{\alpha(1 - \alpha) + \zeta} (1 - g(z)) \right) \quad (2.80)$$

$$\rho_0^{(L)}(q, k) = \frac{3e^2}{2\pi^2} \sum_f Z_f^2 \int_0^1 d\alpha \frac{(\alpha(1 - \alpha))^2}{\alpha(1 - \alpha) + \zeta} (1 - g(z)) \quad (2.81)$$

Here the summation goes over the quark flavours. The dimensionless variables ζ and z are defined as

$$\zeta = \frac{m_f^2}{Q^2}, \quad z = \frac{k^2}{4Q^2} \frac{1}{\alpha(1 - \alpha) + \zeta} \quad (2.82)$$

and m_f and Z_f are the mass and charge of the quark of flavour f . The function $g(z)$ is given by

$$g(z) = \frac{1}{2z\sqrt{z^2 + 1}} \ln \frac{\sqrt{z^2 + 1} + z}{\sqrt{z^2 + 1} - z} \quad (2.83)$$

The structure function is obtained from the cross-section by the standard relation

$$F_2(x, Q^2) = \frac{Q^2}{\pi e^2} (\sigma^{(T)} + \sigma^{(L)}) \quad (2.84)$$

In the DLLA limit only the transverse cross-section contributes. We can also neglect the quark masses in this approximation. Then, with a substitution (2.79), from (2.75), (2.80) and (2.84) we obtain an expression for the quark (sea) density of the target

$$xq(x) = \frac{3}{\pi^2 b_0^2} \frac{Q^2}{\ln Q^2} \int^{Q^2} \frac{dk^2 \phi(x, k)}{k^4 \ln k^2} \int_0^1 d\alpha (\alpha^2 + (1 - \alpha)^2) ((1 + 2z^2)g(z) - 1) \quad (2.85)$$

where $g(z)$ is given by Eq. (2.83) and we assumed that large values of $k^2 < Q^2$ contribute in accordance with the DLLA approximation. In this approximation the asymptotics of the gluon density $xg(x, k^2)$ and consequently of $\phi(x, k^2)$ is known:

$$\phi(x, k^2) = cxg(x, k^2) \simeq c \exp \sqrt{a \ln \frac{1}{x} \ln \ln k^2} \quad (2.86)$$

where $a = 48/b_0$. Putting (2.86) into (2.85), after simple calculations we find the asymptotical expression for the quark density

$$xq(x, k^2) \simeq \frac{4c}{\pi^2 b_0^2} \sqrt{\frac{\ln \ln k^2}{a \ln 1/x}} \exp \sqrt{a \ln \frac{1}{x} \ln \ln k^2} \quad (2.87)$$

On the other hand, let us consider the DGLAP equation in the small x limit (in such a case it is possible to neglect the quark contributions to the sea quarks)

$$\frac{\partial}{\partial \ln Q^2} q(x, Q^2) = \frac{\alpha_s(Q^2)}{8\pi} \int_x^1 \frac{dy}{y} P_{qg}\left(\frac{x}{y}\right) g(y, Q^2) \quad ; \quad P_{qg}(z) = 2[z^2 + (1 - z)^2] \quad (2.88)$$

After a double integration we find

$$xq(x, k^2) \simeq \frac{4}{3b_0^2} \sqrt{\frac{\ln \ln k^2}{a \ln 1/x}} \exp \sqrt{a \ln \frac{1}{x} \ln \ln k^2} \quad (2.89)$$

As we observe the approximation (2.79) for the colour density of the photon projectile leads to the correct relation between the quark and gluon densities in the DLLA limit. This justifies the use of (2.79), at least for high enough $1/x$ and Q^2 . Comparing (2.87) and (2.89) we also obtain the normalization factor c which relates the pomeron wave function to the gluon density

$$c = \pi^2 b_0 / 3 \quad (2.90)$$

2.3.2 The initial distribution

To start the evolution in $1/x$ we have to fix the initial gluon density at some small value $x = x_0$. Evidently, the smaller is x_0 , the smaller is the region where we can compare our predictions

with the experimental data. On the other hand, if x_0 is not small enough, application of the asymptotic hard pomeron theory becomes questionable. Guided by these considerations we choose $x_0 = 0.01$ as our basic initial x although we also tried $x = 0.001$ to see the influence of possible sub-asymptotic effects.

The initial wave function $\phi(x_0, k^2)$ has to be chosen in accordance with the existing data at $x = x_0$ and all k^2 available. The experimental F_2 is a sum of the singlet and non-singlet parts, the latter giving a relatively small contribution at $x = 0.01$. Our theory can give predictions only for the singlet part (and one of the criteria for its applicability is precisely the relative smallness of the non-singlet contribution). The existing experimental data at $x = 0.01$ give values for F_2 averaged over rather large intervals of x and Q^2 . For all these reasons, rather than to try to adjust our initial $\phi(x_0, k^2)$ to the pure experimental data, we have preferred to match it with the theoretical predictions for the gluon density and the singlet part of F_2 given by some standard parametrization fitted to the observed F_2 in a wide interval of Q^2 and small x . As such we have taken the GRV LO parametrization [54]. The choice of LO has been dictated by its comparative simplicity and the fact that at $x = 0.01$ the difference between LO and NLO is insignificant.

Thus, for the initial distribution we have taken the GRV LO gluon density with an appropriate scaling factor. Putting this density into eqs. (2.74), (2.75) and (2.84) one should be able to reproduce the sea quark density and thus the singlet part of the structure function. In the GRV scheme the relation between the gluon density and the quark density is much more complicated and realized through the DGLAP evolution. Since the DGLAP evolution and the pomeron theory are not identical, one should not expect that our initial gluon density should exactly coincide with the GRV one to give the same singlet structure function. One has also to have in mind the approximate character of our colour density ρ at small Q^2 . In fact, with the initial ϕ given by (2.74) and the gluon density exactly taken from the GRV parametrization at $x = 0.01$ we obtain a 30% smaller values for the singlet part of the structure function as given by the same GRV parametrization, the difference growing at low Q^2 . To make the description better we used a certain arbitrariness in the scale Q_1^2 which enters (2.79) and also the scale at which the coupling freezes in the density ρ . A good choice to fit the low Q^2 data is to take

$$\alpha(Q_1^2) = \frac{4\pi}{b_0} \frac{1}{\ln((0.17 * Q^2 + 0.055 (GeV/c)^2)/\Lambda^2)} \quad (2.91)$$

With this $\alpha(Q_1^2)$ the obtained singlet structure function at $x = 0.01$ has practically the same Q^2 dependence as the GRV one, although it results 30% smaller in magnitude. This mismatch can be interpreted in two different ways. Either we may believe that the gluon density given by the GRV is the correct one and the deficiency in the singlet part of the structure function is caused by our approximate form of the colour density ρ (which is most probable). Or we may think that the colour density to be used in the DGLAP should coincide with ours only for large enough Q^2 and $1/x$ and at finite values they may somewhat differ (our relation (2.74) was established strictly speaking only in the DLLA limit). Correspondingly we may either take the relation (2.74) as it stands and use the GRV LO gluon density at $x = 0.01$ in it, or introduce a correcting scaling factor 1.3 which brings the structure function calculated with the help of (2.75)-(2.84) into agreement with the GRV predictions. In the following we adopt the second




Figure 2.11: The singlet part of the structure function of the proton at $x = 0.01$. The continuous line is the result of our calculation while the dashed line correspond to the GRV prediction.

alternative, that is we assume that our initial gluon distribution at $x = 0.01$ is 30% higher than the one given by the GRV parametrization. Of course to strictly compare the rate of growth of the distribution under evolution one should take into account the differences at the starting point.

The singlet part of the structure function at $x = 0.01$ calculated from (2.75)-(2.84) with this choice is shown in Fig. 2.11 together with the GRV predictions. However one can easily pass to the first alternative by simply reducing our results by factor 1.3.

2.3.3 Numerical results

With the initial wave function $\phi(x = 0.01, k^2)$ chosen as indicated in the preceding section we solved the evolution equation for $10^{-8} < x < 10^{-2}$.

The adopted computational scheme consists in diagonalizing the Hamiltonian in (2.70), reduced to one dimension in the transverse momentum space after angular averaging, and represent the initial wave function as a superposition of its eigenvectors. To discretize k^2 a grid was introduced, after which the problem is reduced to a standard matrix one. To check the validity of the obtained results we have also repeated the evolution using a Runge-Kutta method, resulting in a very good agreement. The final results obtained for the gluon distribution $xg(x, Q^2)$ as a function of x for various Q^2 are shown in Fig. 2.12 where the first two plots correspond to x and Q^2 presently available, whereas the last two plots show the behaviour of the calculated gluon density in the region up to very small x and very high Q^2 , well beyond the present possibilities. For comparison we have also shown the gluon densities for the GRV LO parametrization [54], for the MRS parametrization [55] and also for the BFKL evolution as calculated in [56].

Putting the found gluon densities into eqs. (2.75)-(2.84) we obtain the (singlet part of) proton structure function $F_2(x, Q^2)$. The results are illustrated in Fig. 2.13 for the x dependence. As for the gluon densities, the experimentally investigated region is shown in the first two plots, in which the existing experimental data from [10] are also presented.

Finally, to see a possible influence of sub-asymptotic effects, we have repeated the procedure

taking as a starting point for the evolution a lower value $x = 0.001$. The resulting gluon distributions and structure functions are also presented in the above figures.

To discuss the results obtained we have to remember that they involve two quantities of a different theoretical status. One is the pomeron wave function ϕ which can be identified with the gluon distribution (up to a factor). The other is the quark density (which is equivalent to the structure function), for which we actually have no rule for the introduction of a running constant and which in the present calculation involves a semi-phenomenological ansatz (2.79). Evidently the results for the latter are much less informative as to the effect of the running coupling introduced in our way. Therefore we have to separately discuss our prediction for the gluon distribution, on the one hand, and for the structure function, on the other.

Let us begin with the gluon distribution. Comparing our results with those of GRV, which correspond to the standard DGLAP evolution, we observe that at high enough Q^2 and low enough x our distributions rise with Q and $1/x$ faster than those of GRV. This difference is, of course, to be expected. The hard pomeron theory in any version predicts a power rise of the distribution with $1/x$ to be compared with (2.86) for the DGLAP evolution. As to the Q -dependence, the fixed coupling (BFKL) hard pomeron model predicts a linear rise, again much stronger than (2.86). Our running coupling model supposedly leads to a somewhat weaker rise. From our results it follows that it is still much stronger than for the DGLAP evolution. However one can observe that these features of our evolution become clearly visible only at quite high Q and $1/x$. For moderate $Q < 10 \text{ GeV}/c$ and/or $x > 10^{-4}$ the difference between our distributions and those of GRV is insignificant. As to the DGLAP evolved MRS parametrization, it gives the gluon distribution which lies systematically below the GRV one and, correspondingly, below our values, the difference growing with Q and $1/x$.

To compare our gluon evolution dynamics with the BFKL one, as presented in [56], we have to remark that we should take a huge correcting factor that depends in Q^2 for the following reasons: we must remember of the factor 1.3 discussed in section 4 and we should note that the starting values used in [56] for the gluon distribution are lower than the GRV values, by a factor 1.92 at $Q^2 = 4 \text{ GeV}^2/c^2$ and by a factor 1.1 at $Q^2 = 1000 \text{ GeV}^2/c^2$. According to this to compare just the evolution in this two models we should multiply our distribution by a factor that is 0.4 at $Q^2 = 4 \text{ GeV}^2/c^2$ and that is 0.7 at $Q^2 = 1000 \text{ GeV}^2/c^2$. Taking into account this factors looking at Fig. 2.12 we see that there is a good agreement for not too small x values between the two evolutions, while looking at the values at very small x in all the Q^2 interval our model predicts a quite slower growth of the gluon distribution. We note also that the spectral method we have used shows how the dominant components of the spectrum contribute to the evolution of the gluon density; some small changes of the initial gluon distribution may results in different projections onto the spectral basis, but cannot lead to a very different behaviour at very small x , even if, as it is, the initial gluon distribution is well known in a restricted Q^2 interval. This is because the first two dominant states are precisely giving an important contribution in this interval.

Passing to the structure functions we observe in Fig. 2.13 that our results give a somewhat too rapid growth with $1/x$ in the region $10^{-3} < x < 10^{-2}$ as compared to the experimental data (and also to the parametrizations GRV fitted to these data). With the scaling factor 1.3




Figure 2.12: The gluon distributions as a function of x evolved from $x = 0.01$ and $x = 0.001$ for the experimentally accessible kinematical region and for asymptotically high values of Q^2 and $1/x$. Standard DGLAP evolved parametrizations (GRV-LO and MRS) and the BFKL evolved distributions from [56] (we report only few points connected by lines) are shown for comparison. For the latter distribution one should take into account a further factor which actually has been chosen in [56] to reduce the initial distribution (see text).




Figure 2.13: x dependence of the singlet part of the proton structure function obtained by evolution from $x = 0.01$ and $x = 0.001$, compared to the GRV prediction and the ZEUS 94 data. Values for asymptotically high values of Q^2 and $1/x$ are also shown.

introduced to fit the data at $x = 0.01$ we overshoot the data at $x < 10^{-3}$ by $\sim 25\%$. Without this factor we get a very good agreement for $x < 10^{-3}$ but are below experiment at $x = 0.01$ by the same order. This discrepancy may be attributed either to sub-asymptotic effects or to a poor quality of our ansatz (2.79). Comparison with the result obtained with a lower starting point for the evolution $x = 0.001$ shows that sub-asymptotic effects together with a correct form of coupling to quarks may be the final answer.

Chapter 3

Unitarization in the large N_c limit: the three pomeron vertex

In the first chapter we have reviewed and analyzed some aspect and properties of the BFKL approximation to the pomeron which is based on the application of the perturbation theory to QCD in the Regge kinematical region and corresponds to small x_B in processes such as DIS. Specifically the BFKL pomeron is based on the resummation of the leading large logarithms in $1/x$ and presents the important feature of a steep increase of the cross sections as x decreases, with a power like behavior in $1/x$. This is, as we already noted in the first chapter, in contrast with the bound set by unitarity. In fact this is the well known Froissart bound [41], which sets the limit to the increase behavior with energy of the total hadronic cross section to $\ln^2 s$ and which, for DIS, would lead to the growth bound of $\ln^2 1/x$ [42, 43]. Taking the double logarithmic limit, where one resums powers of $(\ln 1/x \ln Q^2)$ and which thus come out also from the small x limit of the DGLAP approach, one has a somewhat less dramatic increase, in accord to an $\exp(\sqrt{1/x})$ behavior, but still unitarity is strongly violated.

Therefore, to extend the kinematical region of validity of the approach one has to go beyond the leading logarithmic approximation. This is in principle a formidable task and could also not be very satisfactory in presence of hadrons which cannot be studied only with perturbative methods. Here we shall be interested only in a perturbative analysis, which nevertheless is better established for the study of processes like onium-onium scattering, and one hopes that any result could give some new insight to the complete theory which surely would need of a non-perturbative treatment.

A systematic approach to take into account a minimal set of corrections necessary to restore the unitarity has been studied by Bartels [19]. The higher order amplitudes, which contain any number of (reggeized) gluons in the t -channel, are constructed using unitarity and dispersion relations starting from the lower order ones and one has a coupled system of equations for the n reggeized gluon state amplitudes. The main feature is the presence of transitions vertices which change the number of reggeon in the t -channel and thus leads to a reggeon field theory.

The study of this problem is far from being complete. The first step has been to consider the coupled equations for the amplitudes relative to 2, 3 and 4 gluons in the t -channel [20]. The most interesting object which appears is an effective vertex for the transition between 2 and 4

gluon in the t -channel. This vertex has been found to be conformal invariant [21] and in the appropriate color subspace is related to the triple pomeron vertex. Another element of some interest is the 4-gluon interacting state (and its generalizations to n gluons) satisfying the so called BKP equation [22, 23]. This amplitude, characterized by a constant number of gluons in the t -channel, has been studied in the large N_c limit and the system has been proved to be complete integrable and equivalent to an XXX Heisenberg zero spin model [24, 25, 26].

The BFKL amplitudes and the $2 \rightarrow 4$ effective reggeon vertex have been interpreted in terms of correlation functions of a 2D conformal field theory [27, 28] hoping to obtain in the future such a formulation for the effective QCD in the Regge limit. In this context some work [57] has been done to understand the spectrum of the 4 gluon state in terms of the scaling dimension of some operators appearing in the twist expansion of the amplitude in the short distance limit. The idea is based on the fact that, due to the strong symmetry of a bidimensional conformal field theory, all the informations are contained in the scaling dimensions of the operators present in the OPE (operator product expansion) and in the expansion coefficients.

We shall be interested here in the study of the coupled system of equations up to 4 reggeized gluons in the t -channel in the large N_c limit [29, 30]. The analysis will be performed at the leading and next to leading order (NLO) in $1/N_c$ since it is at the latter level that the double pomeron amplitude will appear. A link to some results coming from the dipole colour approach let arise the possibility of a general constructions of the n pomeron amplitudes in the large N_c limit. In order to better understand this one should go beyond the 4 gluon system. Already the 6 gluon system in the large N_c limit is capable to show if some equivalence with the A.H. Mueller approach to dipoles is maintained, since it seems that the latter leads to a system of fan diagrams with only a triple pomeron interactions.

We shall start introducing the formal general system of equations for the 4 gluons system and its main properties. Written the system for the large N_c limit case the solutions in the leading order are given and at the NLO in $1/N_c$ system studied. The double pomeron amplitude at NLO is found to separate in a direct double pomeron exchange (DPE) contribution and in a triple pomeron interaction (TPI) term. The TPI term can be written in terms of 3 point function building blocks which are shown to be conformally invariant so that the same property, as already known, has to be shared by the triple pomeron vertex.

It is later shown in detail the interesting fact that the DPE contribution can be substituted by a completely equivalent TPI term (the inverse is not true). The absence of DPE terms was already noted for $N_c = 3$ in [20] and, in fact, in this form the final vertex interactions left essentially coincides in the two approaches, provided one takes the limit $N_c \rightarrow \infty$. Coupled to pomerons, this vertex effectively reduces to the one found by R. Peschanski in [31], in a colour dipole framework. In particular this fact has been contemporaneously also shown by Korchemsky [28]. However this does not mean that the double dipole density in the dipole approach coincides with the diffractive amplitude in the s -channel unitarity approach: there are certain terms in the latter which are missing in the double dipole density and an asymmetry factor which distinguish, in the rapidity evolution, between the original pomeron and the two which originate from the split.

Finally a section is devoted to study some properties of higher order dipole densities in the

A.H. Mueller approach which are found to be represented by a set of pomeron fan diagrams with only a triple pomeron coupling.

3.1 General unitarization approach

A program devoted to considering a minimal subset of next to leading logarithmic contributions necessary to construct a unitary amplitude has been carried out in J. Bartels work [19]. The basic object under consideration is the amplitude D_n for the coupling of n reggeons (gluons) to two external elementary particles in a colour singlet state. These amplitudes result from a multiple discontinuity analysis from which emerges the feature that the number of reggeized gluons in the t -channel is not conserved. This fact can be also analyzed in the physical t -channel region ($t > 0$) looking for the discontinuities along the t -cut associated with the threshold for the reggeon production. One nice feature due to the gluon reggeization property is the absence of contributions, in the adjoint colour representation, to the partial wave amplitude cut related to n -particle production with $n > 2$.

Let us write the formal system of equations in cascade for the D_n amplitudes (seen as vectors in the appropriate colour spaces) up to $n = 4$:

$$\begin{aligned}
\left[j - 1 - \sum_{i=1}^2 \omega(k_i)\right] D_2(\{k_i\}) &= D_{20}(\{k_i\}) + [K_{2 \rightarrow 2} \otimes D_2](\{k_i\}) \\
\left[j - 1 - \sum_{i=1}^3 \omega(k_i)\right] D_3(\{k_i\}) &= D_{30}(\{k_i\}) + [K_{2 \rightarrow 3} \otimes D_2](\{k_i\}) + \sum [K_{2 \rightarrow 2} \otimes D_3](\{k_i\}) \\
\left[j - 1 - \sum_{i=1}^4 \omega(k_i)\right] D_4(\{k_i\}) &= D_{40}(\{k_i\}) + [K_{2 \rightarrow 4} \otimes D_2](\{k_i\}) + \sum [K_{2 \rightarrow 3} \otimes D_3](\{k_i\}) + \\
&\quad \sum [K_{2 \rightarrow 2} \otimes D_4](\{k_i\})
\end{aligned} \tag{3.1}$$

In the above equations D_{n0} is the lowest order term with the gluons coupled directly to the $q\bar{q}$ loop. $K_{2 \rightarrow 2}$ is the BFKL interaction kernel with the gluon propagators included. Remembering the expression for its momentum dependence

$$V(k_1, k_2; k'_1, k'_2) = \frac{k_1^2 k_2'^2 + k_2^2 k_1'^2}{k_1'^2 k_2'^2 (k_1 - k'_1)^2} - \frac{(k_1 + k_2)^2}{k_1'^2 k_2'^2} \tag{3.2}$$

it is possible to give the expression for the kernels $K_{2 \rightarrow 3}$ and $K_{2 \rightarrow 4}$ in a simple way. In fact, defining

$$W(k_1, k_2, k_3; k'_1, k'_3) = V(k_2, k_3; k'_1 - k_1, k'_3) - V(k_1 + k_2, k_3; k'_1, k'_3) \tag{3.3}$$

one has, on including the colour structure,

$$\begin{aligned}
K_{2 \rightarrow 3}(k_1, k_2, k_3; k'_1, k'_3) &= g^3 f^{a'_1 a_1 c} f^{c a_2 d} f^{d a_3 a'_3} W(k_1, k_2, k_3; k'_1, k'_3) \\
K_{2 \rightarrow 4}(k_1, k_2, k_3, k_4; k'_1, k'_4) &= g^4 f^{a'_1 a_1 c} f^{c a_2 d} f^{d a_3 e} f^{e a_4 a'_4} W(k_1, k_2 + k_3, k_4; k'_1, k'_4)
\end{aligned} \tag{3.4}$$

Moreover in the coupled system of equations (3.1) the sums run over all possible pairwise interac-

tions and the symbol $\otimes$ means an integration over primed variables with a weight $(2\pi)^{-3}\delta^{(2)}(\sum k - \sum k')$.

The analysis of the colour structure for the system of 4 gluons was performed by Bartels in [19], separating the gluons in two pairs, each in the same colour quantum state belonging to the set $(1, 8_A, 8_S, \bar{10} + 10, 27)$. Therefore the amplitude has been characterized by a 5-dimensional colour vector for the three different signature assignments $(-, -)$, $(+, +)$, $(+, -)$ and it has been possible to reduce the general operator $K_{4 \rightarrow 4}$, which should appear in principle in the last term of the equation for D_4 , in a sum of $K_{2 \rightarrow 2}$ operators acting on different colour subspaces.

3.2 Large N_c limit for the system of four reggeized gluons

The set of equations (3.1) for the system of four reggeized gluons, coupled to a fermion loop in a colour singlet state, found by J. Bartels and M. Wuesthoff [20] for the real case $N_c = 3$ is quite complicated because of the colour structure.

The large N_c limit [1, 2] is interesting because in the leading order the amplitude for the system is based on a single BFKL pomeron exchange, i.e. one has a fully reggeizing contribution. Performing a perturbative analysis at the NLO in $1/N_c$ one finds a subleading diffractive amplitude which consists of a sum of a DPE and a TPI terms, in full correspondence with the Regge-Gribov picture.

The results presented in this section were studied in [29] and [30] respectively for the forward and for the more general non-forward direction case which is necessary to study also the conformal invariance properties of the $2 \rightarrow 4$ effective gluon vertex.

3.2.1 Leading order in $1/N_c$

The results for a system of two and three gluons are not sensitive to the $N_c \rightarrow \infty$ limit, which instead affects the four gluon case.

Let us first consider the two gluon system. In the lowest (zero) order approximation the 2 gluon amplitude discontinuity (Mellin transformed to the complex angular momentum j) is given by the $q\bar{q}$ loop with the two gluons attached to it in all possible ways (4 diagrams in all). Their colour indices a_1 and a_2 enter into the colour trace $Tr\{t_{a_1}t_{a_2}\} = (1/2)\delta_{a_1a_2}$, where t_a is the colour of the quark. Separating this trace and the coupling g^2 and projecting on the vacuum colour state $|0\rangle = (1/N)\delta_{a_1a_2}$ we write the zero order contribution in the large N_c limit as

$$D_{20}(1, 2) = D_{20}(2, 1) = g^2 N_c (f(1+2, 0) - f(1, 2)) \quad (3.5)$$

where we use the notation in which only the number of the gluon is indicated whose momentum enters as a variable and $f(1, 2) = f(2, 1)$ is a contribution of the $q\bar{q}$ loop with gluon 1 attached to q and gluon 2 attached to $\bar{q}$. Its explicit form can be easily found (see Appendix A.2) but has no importance for the following.

The basic quantity is the full amplitude D_2 . It corresponds to the exchange of two reggeized

gluons (the BFKL amplitude) and satisfies the BFKL equation

$$S_{20}D_2 = D_{20} + g^2 N_c V_{12} D_2 \quad (3.6)$$

where S_{20} is the 2 gluon "free" Schrödinger operator for the energy $1 - j$

$$S_{20} = j - 1 - \omega(1) - \omega(2) \quad (3.7)$$

$\omega(k)$ is, as usual, the gluon Regge trajectory, V_{12} is the BFKL interaction and the previous found D_{20} acts as an inhomogeneous term. For simplicity we leave here the $\otimes$ symbol understood.

For the number of exchanged gluons $n = 3, 4$ amplitudes $D_n(j)$ are defined as integrals of $D_n(j_1, \dots, j_{n-1})$, depending on $n - 1$ partial t -channel angular momenta, over all j_i subject to condition $\sum_{i=1}^{n-1} (j_i - 1) = j - 1$ corresponding to conservation of "reggeon energies". Moreover we extend the notation for the n reggeized gluon free Schrödinger operator defining $S_{n0} = j - 1 - \sum_i \omega(i)$.

The three gluon system is described by an equation which can be solved in terms of the D_2 amplitudes thanks to the gluon reggeization property. It is worth to study its colour structure. The colour factor associated to the lowest order amplitude of three gluons attached to the quark loop is given by the trace $Tr\{t_{a_1} t_{a_2} t_{a_3}\} = (1/4) h_{a_1 a_2 a_3}$, where $h_{a_1 a_2 a_3} = d_{a_1 a_2 a_3} + i f_{a_1 a_2 a_3}$ is cyclic symmetric and satisfies the properties

$$h_{a_1 a_2 a_3}^* = h_{a_2 a_1 a_3} \quad , \quad \sum_{cd} h_{acd}^* h_{bcd} = \delta_{ab} 2N_c (1 - 2/N_c^2) \quad , \quad \sum_{cd} h_{acd} h_{bcd} = -\delta_{ab} (4/N_c) \quad (3.8)$$

One may introduce the two colour wave functions

$$|123\rangle = \frac{1}{\sqrt{2N_c^3}} h_{a_1 a_2 a_3}, \quad |213\rangle = \frac{1}{\sqrt{2N_c^3}} h_{a_2 a_1 a_3} \quad (3.9)$$

orthonormal in the large N_c limit and project the above lowest order amplitude on to these states. The result can be expressed in terms of the function f because the momenta of the gluons attached to the same fermion line sum together; there is also an overall sign depending on the parity of the number of gluons attached to the quark or to the antiquark lines. One has, for $N_c \rightarrow \infty$,

$$D_{30}^{(123)} = -D_{30}^{(213)} = g \sqrt{\frac{N_c}{8}} [D_{20}(2, 1+3) - D_{20}(1, 2+3) - D_{20}(3, 1+2)] \quad (3.10)$$

Due to the global colour singlet state, each pair is in an adjoint representation and the interaction cannot change the colour configuration. Remembering the bootstrap property each two reggeon state in an adjoint representation gives one reggeon state so that the considered three gluon amplitude must reduce to a superposition of BFKL pomeron amplitudes.

In more detail one can write the equation for the colour state $|123\rangle$

$$S_{30} D_3^{(123)} = D_{30}^{(123)} + D_{2 \rightarrow 3}^{(123)} + \frac{g^2 N_c}{2} (V_{12} + V_{23} + V_{31}) D_3 \quad (3.11)$$

The second inhomogeneous term $D_{2 \rightarrow 3}^{(123)}$ is related to the contribution which arises for the transition from 2 to 3 gluons. In fact, using the vertex $K_{2 \rightarrow 3}$ recalled in the previous section and projecting onto the state $|123\rangle$ one finds

$$D_{2 \rightarrow 3}^{(123)} = g^3 \sqrt{\frac{N_c^3}{8}} W(1, 2, 3; 1'3') \otimes D_2(1', 3') \equiv g^3 \sqrt{\frac{N_c^3}{8}} W_2(1, 2, 3) \quad (3.12)$$

In the equation for the orthogonal state $|213\rangle$ the inhomogeneous terms have opposite sign. As expected the solutions of the equations can simply be obtained promoting the D_{20} in the zero order term to a full BFKL pomeron amplitudes D_2 , i.e.

$$D_3^{(123)} = -D_3^{(213)} = g \sqrt{\frac{N_c}{8}} [D_2(2, 1+3) - D_2(1, 2+3) - D_2(3, 1+2)] \quad (3.13)$$

This can be easily proved (see appendix A.4) directly using (3.3) to transform $D_{2 \rightarrow 3}^{(123)}$ and by applying the bootstrap condition

$$\frac{g^2 N_c}{2} \int \frac{d^2 k'_1}{(2\pi)^3} V(k_1, k_2; k'_1, k'_2) \equiv \frac{g^2 N_c}{2} V_{12} \otimes 1 = \omega(1) + \omega(2) - \omega(1+2) \quad (3.14)$$

to obtain contributions coming from BFKL equations for amplitudes where two reggeized gluons have fused into one.

Let us now consider the four gluon system. In the leading order the colour configuration is much more conveniently described by the order of gluons along the cylinder surface 1234, 1324, etc. If one takes the lowest order amplitude given by the contributions of 16 diagrams with the 4 gluons attached in all possible ways to the $q\bar{q}$ loop, one has for each case a color factor given by the trace $Tr\{t_{a_1} t_{a_2} t_{a_3} t_{a_4}\} = (1/8)h_{a_1 a_2 b} h_{a_3 a_4 b} + (1/4 N_c) \delta_{a_1 a_2} \delta_{a_3 a_4}$. In the $N_c \rightarrow \infty$ limit only the first term, cyclic symmetric, survives. It is useful to define the colour state

$$|1234\rangle = \frac{1}{2N_c^2} h_{a_1 a_2 b} h_{a_3 a_4 b} \quad (3.15)$$

together with others, differing for a permutation, orthonormalized in the large N_c limit. The zero order amplitude projected on to these states gives the components

$$D_{40}^{(1234)} = D_{40}^{(4321)} = \frac{g^2 N_c}{4} [D_{20}(1, 2+3+4) + D_{20}(4, 1+2+3) - D_{20}(1+4, 2+3)] \quad (3.16)$$

$$D_{40}^{(2134)} = D_{40}^{(4312)} = \frac{g^2 N_c}{4} [D_{20}(2, 1+3+4) + D_{20}(3, 1+2+4) - D_{20}(1+2, 3+4) - D_{20}(1+3, 2+4)] \quad (3.17)$$

The gluons lie on a cylinder attached to the $q\bar{q}$ loop and all the neighbouring pairs are globally in an adjoint representation. Moreover in the leading order in $1/N_c$ all the interactions across the cylinder are suppressed and only the neighbouring gluons interact.

The equations for the above amplitudes decouple and are given by [29]

$$\begin{aligned} S_{40}D_4^{(1234)} &= D_{40}^{(1234)} + D_{2 \rightarrow 4}^{(1234)} + D_{3 \rightarrow 4}^{(1234)} + \frac{g^2 N_c}{2}(V_{12} + V_{23} + V_{34} + V_{41})D_4^{(1234)} \\ S_{40}D_4^{(2134)} &= D_{40}^{(2134)} + D_{3 \rightarrow 4}^{(2134)} + \frac{g^2 N_c}{2}(V_{21} + V_{13} + V_{34} + V_{42})D_4^{(2134)} \end{aligned} \quad (3.18)$$

The inhomogeneous terms related to the transitions which change the number of gluons from 2 to 4 are

$$D_{2 \rightarrow 4}^{(1234)} = -\frac{g^4 N_c^2}{4}W_2(1, 2 + 3, 4) \quad (3.19)$$

while the transitions between 3 and 4 gluons are due to the terms

$$\begin{aligned} D_{3 \rightarrow 4}^{(1234)} &= g^3 \sqrt{\frac{N_c^3}{8}} \left[W(2, 3, 4; 2', 4') \otimes D_3^{(124)}(1, 2', 4') + W(1, 2, 3; 1', 3') \otimes D_3^{(134)}(1', 3', 4) \right] \\ D_{3 \rightarrow 4}^{(2134)} &= -g^3 \sqrt{\frac{N_c^3}{8}} \left[W(1, 2, 4; 1', 4') \otimes D_3^{(134)}(1', 3, 4') + W(1, 3, 4; 1', 4') \otimes D_3^{(124)}(1', 2, 4') \right] \end{aligned} \quad (3.20)$$

It can be shown that also the solutions for the 4 gluon amplitudes can be written in terms of the two reggeon amplitudes just by substituting the D_{20} terms in (3.16) and (3.17) by D_2 ones. Thus in the leading order in $1/N_c$ the 4 reggeized gluon system reduces to a single pomeron.

3.2.2 Next to leading order in $1/N_c$

The corrections at the NLO in $1/N_c$ force to consider the two subsystems of gluons (12) and (34). The state which relates to the diffractive amplitude and contributes to the triple pomeron interaction is characterized by both the subsystems in a colour singlet state so that we shall project the zero order state of the 4 gluons attached to the quark loop onto $|0\rangle = (1/N^2)\delta_{a_1 a_2} \delta_{a_3 a_4}$, obtaining

$$D_{40}^{(0)} = \frac{1}{2}g^2 \left(\sum_{i=1}^4 D_{20}(i, 1 + 2 + 3 + 4 - i) - \sum_{i=2}^4 D_{20}(1 + i, 2 + 3 + 4 - i) \right) \quad (3.21)$$

which is explicitly down by a factor $1/N_c$ respect to the 4 gluons amplitudes at leading order (its order is $g^4 N_c$ compared to $g^4 N_c^2$ of the amplitudes $D_{40}^{(1234)}$ and $D_{40}^{(2134)}$). The full diffractive amplitude $D_4^{(0)}$ (directly related to the diffractive cross-section integrated over the diffractive mass) results to be the solution of the following equation

$$S_{40}D_4^{(0)} = D_{40}^{(0)} + D_{2 \rightarrow 4}^{(0)} + D_{3 \rightarrow 4}^{(0)} + D_{4 \rightarrow 4}^{(0)} + g^2 N_c (V_{12} + V_{34})D_4^{(0)} \quad (3.22)$$

where there is also an inhomogeneous term $D_{4 \rightarrow 4}^{(0)}$ which comes from a LO 4 gluon cylindric configuration due to the interactions between adjacent gluons as well as across the cylinder. In the Bartels approach [19] the corresponding transitions are related to the matrix elements 12 and 13 (which turns out to be zero) of the matrix $K_{4 \rightarrow 4}^T$. The contributions to the two colour

singlet amplitude at the NLO have explicitly been found [29]:

$$\begin{aligned}
D_{2 \rightarrow 4}^{(0)} &= -g^4 N_c W_2(1, 2 + 3, 4) \\
D_{3 \rightarrow 4}^{(0)} &= g^3 \sqrt{2N_c} \left[W(1, 2, 3; 1'3') \otimes D_3^{(134)}(1', 3', 4) - W(1, 2, 4; 1', 4') \otimes D_3^{(134)}(1', 3, 4') + \right. \\
&\quad \left. W(2, 3, 4; 2', 4') \otimes D_3^{(124)}(1, 2', 4) - W(1, 3, 4; 1', 4') \otimes D_3^{(124)}(1', 2, 4') \right] \\
D_{4 \rightarrow 4}^{(0)} &= g^2 (V_{23} + V_{14} - V_{13} - V_{24}) \left[D_4^{(1234)} - D_4^{(2134)} \right]
\end{aligned} \tag{3.23}$$

From the inspection of these inhomogeneous terms one can observe that each one is given by the action of some operator on the BFKL pomeron amplitude D_2 . It is then possible to define an operator Z associated to the triple pomeron vertex and acting on the BFKL pomeron

$$Z D_2 \equiv D_{2 \rightarrow 4}^{(0)} + D_{3 \rightarrow 4}^{(0)} + D_{4 \rightarrow 4}^{(0)} \tag{3.24}$$

and rewrite eq. (3.22) as

$$S_{40} D_4^{(0)} = D_{40}^{(0)} + Z D_2 + g^2 N_c (V_{12} + V_{34}) D_4^{(0)} \tag{3.25}$$

The three-pomeron vertex operator Z can be conveniently reexpressed in terms of a new function $G(k_1, k_2, k_3)$ (see also appendix A.3) which is defined as the vertex $K_{2 \rightarrow 3}$ for the transition of 2 to 3 gluons, integrated with the BFKL pomeron and regularized in the infrared by terms proportional to the gluon trajectory in the same manner as in the total BFKL kernel

$$\begin{aligned}
G(k_1, k_2, k_3) &= G(k_3, k_2, k_1) = -g^2 N_c W_2(k_1, k_2, k_3) - D(k_1, k_2 + k_3) (\omega(k_2) - \omega(k_2 + k_3)) - \\
&\quad D(k_1 + k_2, k_3) (\omega(k_2) - \omega(k_1 + k_2))
\end{aligned} \tag{3.26}$$

It is a generalization of a similar function of two momenta introduced in [20] for the forward case and in the graphical notation introduced by Bartels it can be written as illustrated in Fig. 3.1.

If we write $D_{3 \rightarrow 4}^{(0)}$ using G we obtain (as usual we denote k_i by writing just i for brevity),

$$\begin{aligned}
D_{3 \rightarrow 4}^{(0)} &= \frac{g^2}{2} \left[2 G(2, 3 + 4, 1) + 2 G(3, 1 + 2, 4) - G(1 + 3, 2, 4) - G(1 + 4, 2, 3) - \right. \\
&\quad G(1 + 4, 3, 2) - G(2 + 4, 3, 1) + \\
&\quad D_2(1, 2 + 3 + 4) (2\omega(3 + 4) - \omega(3) - \omega(2 + 3 + 4)) + \\
&\quad D_2(2, 1 + 3 + 4) (2\omega(3 + 4) - \omega(3) - \omega(1 + 3 + 4)) + \\
&\quad D_2(3, 1 + 2 + 4) (2\omega(1 + 2) - \omega(2) - \omega(1 + 2 + 4)) + \\
&\quad D_2(4, 1 + 2 + 3) (2\omega(1 + 2) - \omega(2) - \omega(1 + 2 + 3)) + \\
&\quad D_2(1 + 3, 2 + 4) (\omega(2 + 4) - \omega(2) - \omega(3)) + \\
&\quad \left. D_2(1 + 4, 2 + 3) (\omega(2 + 3) - \omega(2) - \omega(3)) \right]
\end{aligned} \tag{3.27}$$

This expression is not symmetric in gluons (12) nor in gluons (34) and, since the function $D_4^{(0)}$ is symmetric in both gluon pairs, one should symmetrize it. The resulting symmetric expression




Figure 3.1: Graphical representation of the conformal invariant function G in term of which the vertex Z is constructed: for each line one has to write the corresponding propagator, for each vertex the square of the sum of the momenta above or below that vertex and the propagators of the lower lines are omitted.

is

$$\begin{aligned}
D_{3 \rightarrow 4}^{(0)} = & \frac{g^2}{4} \left[2 G(1, 3+4, 2) + 2 G(2, 3+4, 1) + 2 G(3, 1+2, 4) + 2 G(4, 1+2, 3) - \right. \\
& G(1+3, 2, 4) - G(1+3, 4, 2) - G(2+3, 1, 4) - G(2+3, 4, 1) - \\
& G(1+4, 2, 3) - G(1+4, 3, 2) - G(2+4, 1, 3) - G(2+4, 3, 1) + \\
& D_2(1, 2+3+4) (2 \omega(2+3+4) - 2 \omega(2+4) - 2 \omega(2+3) + \omega(3) + \omega(4)) + \\
& D_2(2, 1+3+4) (2 \omega(1+3+4) - 2 \omega(1+4) - 2 \omega(1+3) + \omega(3) + \omega(4)) + \\
& D_2(3, 1+2+4) (2 \omega(1+2+4) - 2 \omega(2+4) - 2 \omega(1+4) + \omega(1) + \omega(2)) + \\
& D_2(4, 1+2+3) (2 \omega(1+2+3) - 2 \omega(1+3) - 2 \omega(2+3) + \omega(1) + \omega(2)) \\
& D_2(1+3, 2+4) (2 \omega(1+3) + 2 \omega(2+4) - \omega(1) - \omega(2) - \omega(3) - \omega(4)) + \\
& \left. D_2(1+4, 2+3) (2 \omega(1+4) + 2 \omega(2+3) - \omega(1) - \omega(2) - \omega(3) - \omega(4)) \right] \quad (3.28)
\end{aligned}$$

The term $D_{2 \rightarrow 4}^{(0)}$ can be rewritten as

$$D_{2 \rightarrow 4}^{(0)} = g^2 \left[G(1, 2+3, 4) + D_2(1, 2+3+4)(\omega(2+3) - \omega(2+3+4)) + D_2(4, 1+2+3)(\omega(2+3) - \omega(1+2+3)) \right] \quad (3.29)$$

and, after an analogous symmetrization procedure, it becomes

$$\begin{aligned}
D_{2 \rightarrow 4}^{(0)} = & (1/4) g^2 \left[G(1, 2+3, 4) + G(1, 2+4, 3) + G(2, 1+3, 4) + G(2, 1+4, 3) + \right. \\
& D_2(1, 2+3+4)(\omega(2+3) + \omega(2+4) - 2 \omega(2+3+4)) + \\
& D_2(2, 1+3+4)(\omega(1+3) + \omega(1+4) - 2 \omega(1+3+4)) + \\
& \left. D_2(3, 1+2+4)(\omega(1+4) + \omega(2+4) - 2 \omega(1+2+4)) + \right.
\end{aligned}$$

$$D_2(4, 1 + 2 + 3)(\omega(1 + 3) + \omega(2 + 3) - 2 \omega(1 + 2 + 3)) \Big] \quad (3.30)$$

We finally come to the term $D_{4 \rightarrow 4}^{(0)}$. The operator $V_{23} + V_{14} - V_{13} - V_{24}$ is antisymmetric under the interchange of 1 and 2 and/or 3 and 4. Since the two-pomeron state is symmetric under these substitutions, only antisymmetric parts of $D_4^{(1234)}$ and $D_4^{(2134)}$ give a nonzero contribution. Taking this into account we find

$$D_{4 \rightarrow 4}^{(0)} = \frac{g^4 N_c}{4} (V_{23} + V_{14} - V_{13} - V_{24}) [D_2(1 + 3, 2 + 4) - D_2(1 + 4, 2 + 3)] \quad (3.31)$$

As we have previously noted these terms are due to the interaction between different gluons (e.g. $V_{13}D_2(1 + 4)$) or within the same gluon (e.g. $V_{13}D_2(1 + 3)$).

The first type gives the contribution

$$D_{4 \rightarrow 4}^{(01)} = \frac{g^4 N_c}{4} [(V_{13} + V_{24})D_2(1 + 4) + (V_{23} + V_{14})D_2(1 + 3)] \quad (3.32)$$

To reduce it to the expression of the same type as before we use the identity (see appendix A.4):

$$V_{13}D_2(1 + 4) = W_2(2 + 3, 1, 4) + W_2(2, 3, 1 + 4) - W_2(2, 1 + 3, 4) - W_2(2 + 3, 0, 1 + 4) \quad (3.33)$$

Introducing then functions G we find for this part

$$\begin{aligned} D_{4 \rightarrow 4}^{(01)} = & \frac{g^4 N_c}{4} \Big[G(1, 2 + 3, 4) + G(1, 2 + 4, 3) + G(2, 1 + 3, 4) + G(2, 1 + 4, 3) - \\ & G(1, 3, 2 + 4) - G(1, 4, 2 + 3) - G(2, 3, 1 + 4) - G(2, 4, 1 + 3) - \\ & G(3, 1, 2 + 4) - G(3, 2, 1 + 4) - G(4, 1, 2 + 3) - G(4, 2, 1 + 3) + \\ & 2 G(1 + 3, 0, 2 + 4) + 2 G(1 + 4, 0, 2 + 3) + \\ & D_2(1, 2 + 3 + 4)(\omega(2 + 3) + \omega(2 + 4) - \omega(3) - \omega(4)) + \\ & D_2(2, 1 + 3 + 4)(\omega(1 + 3) + \omega(1 + 4) - \omega(3) - \omega(4)) + \\ & D_2(3, 1 + 2 + 4)(\omega(2 + 4) + \omega(1 + 4) - \omega(1) - \omega(2)) + \\ & D_2(4, 1 + 2 + 3)(\omega(1 + 3) + \omega(2 + 3) - \omega(3) - \omega(4)) - \\ & (D_2(1 + 3, 2 + 4) + D_2(1 + 4, 2 + 3)) \sum_{i=1}^4 \omega(i) \Big] \quad (3.34) \end{aligned}$$

The second part leads to the contribution

$$\begin{aligned} D_{4 \rightarrow 4}^{(02)} = & -\frac{g^4 N_c}{4} [(V_{13} + V_{24})D_2(1 + 3, 2 + 4) + \\ & (V_{23} + V_{14})D_2(1 + 4, 2 + 3)] \quad (3.35) \end{aligned}$$

Applying the bootstrap relation one immediately has

$$D_{4 \rightarrow 4}^{(02)} = \frac{g^2 N_c}{2} \left[D_2(1+3, 2+4) \left(\sum_{i=1}^4 \omega(i) - \omega(1+4) \right) - \omega(2+3) + \right. \\ \left. D_2(1+3, 2+4) \left(\sum_{i=1}^4 \omega(i) - \omega(1+3) - \omega(2+4) \right) \right] \quad (3.36)$$

The final three-pomeron vertex is obtained after summing all the contributions, explicitly $ZD_2 = D_{2 \rightarrow 4}^{(0)} + D_{3 \rightarrow 4}^{(0)} + D_{4 \rightarrow 4}^{(01)} + D_{4 \rightarrow 4}^{(02)}$ and is given by:

$$ZD_2 = -\frac{g^2}{2} \left[G(1+3, 2, 4) + G(1+3, 4, 2) + G(1, 3, 2+4) + G(3, 1, 2+4) + \right. \\ G(1+4, 2, 3) + G(1+4, 3, 2) + G(1, 4, 2+3) + G(4, 1, 2+3) - \\ 2G(1, 3+4, 2) - 2G(3, 1+2, 4) - G(1, 2+3, 4) - G(2, 1+4, 3) - \\ \left. G(1, 2+4, 3) - G(2, 1+3, 4) - G(1+3, 0, 2+4) - G(1+4, 0, 2+3) \right] \quad (3.37)$$

One may note from the form of the $K_{2 \rightarrow 3}$ vertex that, after multiplying by the gluon propagators, the G function vanishes if the first or the third argument is zero, but the property is not true for the second argument. However at the global vertex level, after having performed the sum (3.37) of all the contributions, the validity of the Ward identities is restored.

Let us end this section looking at the form of the solution of the equation (3.25). In fact it can be easily solved and, evidently, the solution may be constructed as a sum of two terms corresponding to the two parts of the inhomogeneous term $D_{40}^{(0)}$ and ZD_2 :

$$D_4^{(0)} = D_4^{DPE} + D_4^{TPI} \quad (3.38)$$

The DPE term D_4^{DPE} comes from the inhomogeneous term $D_{40}^{(0)}$ and its explicit form can be conveniently written using the quark loop density in the transverse coordinate space defined by the Fourier transform (see Appendix A.2 for explicit expressions)

$$f(1, 2) = \int d^2 r \rho_l(r) e^{ik_1 r} \quad (3.39)$$

where $l = k_1 + k_2$ and r is the dipole transverse dimension. Thus one has a contribution of the form

$$D_4^{DPE} = (1/4) g^4 N_c \int d^2 r \rho_l(r) D_4^{(r)} \quad (3.40)$$

Here $D_4^{(r)}$ is a convolution in the "energy" $1-j$ of two independent BFKL pomerons

$$D_4^{(r)} = \int dj_{12} dj_{34} \delta(j+1-j_{12}-j_{34}) D_{2,j_{12}}^{(r)}(1, 2) D_{2,j_{34}}^{(r)}(3, 4) \quad (3.41)$$

and to restore the structure of the inhomogeneous term $D_4^{(0)}$ the amplitudes must belong to two

equations with a proper inhomogeneous part. $D_{2,j}^{(r)}(1, 2)$ satisfies the equation

$$S_{20}D_{2,j}^{(r)} = \prod_{j=1}^2 (e^{ik_j r} - 1) + g^2 N_c V_{12} D_{2,j}^{(r)} \quad (3.42)$$

and a similar relation regards the second pomeron. In the rapidity space one can directly write this contribution as a product of two BFKL amplitudes relative to the two colour singlet pairs

$$D_4^{(r)}(y) = D_2^{(r,12)}(y) D_2^{(r,34)}(y) \quad (3.43)$$

The part D_4^{TPI} is the TPI contribution. It can be written as a convolution in the rapidity space:

$$D_4^{TPI}(1, 2, 3, 4; Y) = \int_0^Y G_2(1, 2; 1'2'; Y - y) G_2(3, 4; 3'4'; Y - y) \otimes Z(1', 2', 3', 4'; 1'', 2'') \otimes D_2(1'', 2''; y) \quad (3.44)$$

where G_2 is the BFKL Green function and the symbols $\otimes$ mean integrations over intermediate momenta. This equation clearly shows that Z is just the three-pomeron vertex.

3.3 Conformal invariance

In [20] it was introduced the vertex $V(1234)$. For a non-zero momentum flow in the t -channel it can be written in a way similar to (3.37),

$$V(1234)D_2 = \frac{1}{2}g^2 \left[G(1, 2 + 3, 4) + G(2, 1 + 3, 4) + G(1, 2 + 4, 3) + G(2, 1 + 4, 3) \right. \\ \left. - G(1 + 2, 3, 4) - G(1 + 2, 4, 3) - G(1, 2, 3 + 4) - G(2, 1, 3 + 4) + G(1 + 2, 0, 3 + 4) \right] \quad (3.45)$$

This vertex is conformal invariant [21] in the following sense. If one transforms VD_2 to the transverse coordinate space and integrates it over the 4 gluon coordinates with a conformal invariant function, the resulting integral is invariant under conformal transformation of gluon coordinates. Comparing (3.45) and (3.37) one immediately see that our vertex Z can be written as

$$Z = V(1324) + V(1423) \quad (3.46)$$

and thus is conformal invariant.

However a stronger conformal property is satisfied: not only the combination (3.45) of functions G is conformal invariant, but each function $G(1, 2, 3)$ is conformal invariant by itself. This is related to the fact that the function G defined in (3.26) represents a natural generalization of the BFKL kernel not only in respect to its infrared stability but also in its conformal properties.

To prove the previous statement one needs a representation of G in the coordinate space. We give all the technical details in appendix A.3 and use here the computed final expressions.

Denote the integral part of G , given by the W_2 term in (3.26) as G_1 and the remaining terms

related to the gluon trajectories as G_2 . In A.3 the following expression is derived:

$$G_1(r_1, r_2, r_3) = A_1 D_2(r_1, r_3) \quad (3.47)$$

where A_1 is an operator in the coordinate space

$$A_1 = \frac{g^2 N_c}{8\pi^3} \left[2\pi\delta^2(r_{23})\partial_3^2(c - \ln r_{13})\partial_3^{-2} + 2\pi\delta^2(r_{12})\partial_1^2(c - \ln r_{13})\partial_1^{-2} \right. \\ \left. - 2\frac{\mathbf{r}_{12}\mathbf{r}_{23}}{r_{12}^2 r_{23}^2} - 2\pi(c - \ln r_{13})(\delta^2(r_{12}) + \delta^2(r_{23})) - 4\pi^2\delta^2(r_{12})\delta^2(r_{23})(\partial_1 + \partial_3)^2\partial_1^{-2}\partial_3^{-2} \right] \quad (3.48)$$

Here $r_{ij} = r_i - r_j$ and $c = \ln(2/m) + \psi(1)$ with m the gluon mass acting as an infrared cutoff.

The transformation of the part G_2 requires introduction of an ultraviolet cutoff ϵ due to the presence of the gluon trajectory terms. Of course the final results do not depend on ϵ . One obtains

$$G_2(r_1, r_2, r_3) = A_2 D_2(r_1, r_3) \quad (3.49)$$

where A_2 is another operator in the coordinate space

$$A_2 = -\frac{g^2 N_c}{8\pi^3} \left[\frac{1}{r_{23}^2} - 2\pi c\delta^2(r_{23}) \right] + \delta^2(r_{23})\omega(-i\partial_3) - \frac{g^2 N_c}{8\pi^3} \left[\frac{1}{r_{12}^2} - 2\pi c\delta^2(r_{12}) \right] + \delta^2(r_{12})\omega(-i\partial_1) \quad (3.50)$$

The four terms in A_2 correspond to the four respective terms in G_2 . We remember that the singular operators $1/r_{12}^2$ and $1/r_{23}^2$ are in fact defined with the help of ϵ as

$$\frac{1}{r^2} \equiv \frac{1}{r^2 + \epsilon^2} + 2\pi\delta^2(r)\ln\epsilon, \quad \epsilon \rightarrow 0 \quad (3.51)$$

which is the expression in (A.69) and does not depend on ϵ . Summing A_1 and A_2 the terms containing $\ln m$ cancel, thus the dependence on the gluon mass disappears and $G(r_1, r_2, r_3)$ is infrared stable.

We want to check the conformal invariance of the integral

$$I = \int d^2 r_1 d^2 r_2 d^2 r_3 \Phi(r_1, r_2, r_3) G(r_1, r_2, r_3) \quad (3.52)$$

where function Φ is conformal invariant. We shall demonstrate that the integral I does not change under conformal transformations. In doing so we shall use the fact that the BFKL solution $\Psi(r_1, r_2) = \partial_1^{-2}\partial_2^{-2}D_2(r_1, r_2)$ is conformal invariant, as it has been shown in the first chapter.

It is sufficient to study the behaviour of the function G only under the inversion since the invariance under translations and rescaling is obvious. The complex notation (see section 3 of the first chapter) is more convenient for this aim: under inversion one has

$$r \rightarrow 1/r, \quad k \equiv -i\partial \rightarrow r^2 k \quad (3.53)$$

Therefore (for real r)

$$d^2r \rightarrow d^2r/r^4 \quad (3.54)$$

and

$$D_2(r_1, r_2) \rightarrow r_1^4 r_2^4 D_2(r_1, r_2) \quad (3.55)$$

There are some contributions which are evidently invariant under inversion. For example taking the last term from A_1 it is easy to check that it leads to the integral

$$I_1 = \frac{g^2 N_c}{2\pi} \int d^2r \Phi(r, r, r) \partial^2 \Psi(r, r) \quad (3.56)$$

which is conformal invariant, since both functions Φ and Ψ are invariant, and the factor r^{-4} from d^2r is cancelled by the factor r^4 from ∂^2 .

Terms with the denominators r_{12}^2 and/or r_{23}^2 from $A_1 + A_2$ combine into an integral

$$I_2 = -\frac{g^2 N_c}{8\pi^3} \int d^2r_1 d^2r_2 d^2r_3 \Phi(r_1, r_2, r_3) \frac{r_{13}^2}{r_{12}^2 r_{23}^2} D_2(r_1, r_3) \quad (3.57)$$

in which the regularization (3.51) is implied. Under inversion the ultraviolet cutoff ϵ is transformed into $\epsilon_1 = r_1 r_2 \epsilon$ and $\epsilon_2 = r_2 r_3 \epsilon$ in the denominators r_{12}^2 and r_{23}^2 respectively. Consequently one finds a change of I_2 :

$$\Delta I_2 = \frac{g^2 N_c}{4\pi^2} \int d^2r_1 d^2r_3 \left[\Phi(r_1, r_1, r_3) \ln r_1^2 + \Phi(r_1, r_3, r_3) \ln r_3^2 \right] D_2(r_1, r_3) \quad (3.58)$$

All the other terms in $A_1 + A_2$, proportional either to $\delta^2(r_{12})$ or to $\delta^2(r_{23})$ can be divided into two parts. On one side one has terms in which the δ -function is multiplied either by a constant or by $\ln r_{13}$. It will give rise to a part of the integral I_3 . Terms with a constant are evidently invariant under inversion. However those containing $\ln r_{13}$ are not and lead to the corresponding change of I_3

$$\Delta I_3 = -\frac{g^2 N_c}{8\pi^2} \int d^2r_1 d^2r_3 \left[\Phi(r_1, r_3, r_3) + \Phi(r_1, r_1, r_3) \right] \ln(r_1^2 r_3^2) D_2(r_1, r_3) \quad (3.59)$$

The second part contains differential operators acting on D_2 of an Hamiltonian type. It has a form

$$-\frac{g^2 N_c}{8\pi^2} \left[a_1 \delta^2(r_{12}) + a_3 \delta^2(r_{23}) \right] \quad (3.60)$$

where

$$a_1 = \partial_1^2 \ln r_{13}^2 \partial_1^{-2} + \ln(-\partial_1^2) \equiv \partial_1^2 \tilde{a}_1 \partial_1^{-2} \quad (3.61)$$

and a_3 is obtained by interchange $1 \leftrightarrow 3$. As shown in the section 1.3 the operator a_1 can be transformed into a different form more convenient to analyze its properties under inversion. Indeed, in the complex notation, one writes the expression:

$$\tilde{a}_1 = \ln r_{13} + \ln k_1 + c.c \quad (3.62)$$

A useful relation, between pseudo-differential operators,

$$\ln r_{13} + \ln k_1 = \ln(r_{13}^2 k_1) - k_1^{-1} \ln r_{13} k_1 \quad (3.63)$$

can be deducted from the eq. (1.43). On applying it one finds

$$a_1 = k_1 \ln(r_{13}^2 k_1) k_1^{-1} - \ln r_{13} + c.c. \quad (3.64)$$

Since under the inversion we have

$$\ln r_{13}^2 k_1 \rightarrow \ln r_{13}^2 k_1 - 2 \ln r_3 \quad (3.65)$$

the change in a_1 is

$$\Delta a_1 = -2 \ln r_{13} + \ln(r_1 r_3) + c.c. = \ln \frac{r_1^2}{r_3^2} \quad (3.66)$$

and the variation of the last part of the integral I_4 , which comes from (3.60), is

$$\Delta I_4 = -\frac{g^2 N}{8\pi^2} \int d^2 r_1 d^2 r_3 \left[\Phi(r_1, r_1, r_3) - \Phi(r_1, r_3, r_3) \right] D_2(r_1, r_3) \ln \frac{r_1^2}{r_3^2} \quad (3.67)$$

Summing all the changes one has $\Delta I_2 + \Delta I_3 + \Delta I_4 = 0$ and this proves that the total integral I is indeed invariant under inversion and, hence, under a global conformal transformation.

3.4 Double pomeron exchange and triple pomeron interaction

We already know the general form of the solution of the 4-gluon equation for the diffractive amplitude

$$S_{40} D_4^{(0)} = D_{40}^{(0)} + Z D_2 + g^2 N_c (V_{12} + V_{34}) D_4^{(0)} \quad (3.68)$$

It can be constructed as a sum of the DPE and a TPI parts as in eq. (3.38) from the sum of various inhomogeneous terms.

However a more detailed analysis is instructive. Let us separate from this known exact solution some arbitrary function f , which may depend on the angular momentum j :

$$D_4^{(0)}(j) = f(j) + \tilde{D}_4^{(0)}(j) \quad (3.69)$$

This choice leads to an equation for the new 4-gluon function $\tilde{D}_4^{(0)}$

$$S_{40} \tilde{D}_4^{(0)} = D_{40}^{(0)} + Z D_2 - [S_{40} - g^2 N_c (V_{12} + V_{34})] f + g^2 N_c (V_{12} + V_{34}) \tilde{D}_4^{(0)} \quad (3.70)$$

where the total inhomogeneous part has changed

$$D_{40}^{(0)} + Z D_2 \rightarrow D_{40}^{(0)} + Z D_2 - X \quad , \quad X = [S_{40} - g^2 N_c (V_{12} + V_{34})] f \quad (3.71)$$

This procedure seems quite trivial but, as we shall see, gives some insight into the structure of the eq. (3.68) and of its solution. To this end we choose f to be the BFKL function

which depends on some gluon momenta. The idea of separating a reggeization piece which correspond to BFKL pomerons has already been used in [20] and we shall try here to see in detail how it works; in this way we shall also be able to compare the results obtained in the two approaches. The separation of this simple term f in the amplitude leads to the subtraction of a new inhomogeneous term X which acquires the structure of a triple pomeron term. This means also that one can calculate some specific triple pomeron contributions expressing them in terms of simple functions.

Let us see how this procedure works in some important cases, differing for the momenta which enter in the separated BFKL function. We shall find that, with a particular simple choice for f , one is able to completely cancel the $D_{40}^{(0)}$ which generates the DPE contribution. In the following, for simplicity, only the forward case is considered.

Let $f = (1/2)g^2 D_2(1) = (1/2)g^2 D_2(2 + 3 + 4)$. One has

$$X = \frac{g^2}{2} [S_{40} - g^2 N_c (V_{12} + V_{34})] D_2(1) \quad (3.72)$$

Using eq. (3.6) we can express the $j - 1$ term in S_{40} in terms of the forward BFKL interaction V_0 and ω and have

$$X = \frac{g^2}{2} D_{20} + \frac{g^2}{2} [g^2 N_c (V_0 - V_{12} - V_{34}) + 2\omega(1) - \sum \omega(i)] D_2(1) \quad (3.73)$$

Making use of the relation

$$(V_0 - V_{12}) D_2(1) = -W_2(1, 2, 3 + 4) \quad (3.74)$$

where W_2 is defined by (3.12) and (3.3), applying the bootstrap for the (34) gluon system

$$g^2 N_c V_{34} D_2(1) = 2(\omega(3 + 4) - \omega(3) - \omega(4)) D_2(1) \quad (3.75)$$

and introducing the function $G(1, 3)$ with two arguments [20], which coincides with $G(1, 2, 3)$ for the forward direction case ($1 + 2 + 3 = 0$), we finally obtain

$$X = \frac{g^2}{2} [D_{20}(1) + G(1, 3 + 4) + D_2(1)(\omega(3) + \omega(4) - 2\omega(3 + 4)) + D_2(1 + 2)(\omega(2) - \omega(1 + 2))] \quad (3.76)$$

Thus we find that the changed function $\tilde{D}_4^{(0)}$ will satisfy an equation with a new inhomogeneous term

$$D_{40} + Z D_2 + \frac{g^2}{2} [-D_{20}(1) - G(1, 3 + 4) - D_2(1)(\omega(3) + \omega(4) - 2\omega(3 + 4)) - D_2(1 + 2)(\omega(2) - \omega(1 + 2))] \quad (3.77)$$

Note that the additional term $-(1/2)g^2 D_{20}$ will cancel the identical term in $D_{40}^{(0)}$. As a result, one has converted the double pomeron exchange contribution coming from $(1/2)g^2 D_{20}(1)$ into a triple pomeron contribution corresponding essentially to $G(1, 3 + 4)$ and in the same time has explicitly separated the term $(1/2)g^2 D_2(1)$ in the amplitude. In other words, one can calculate

the triple pomeron contribution corresponding to a vertex

$$\frac{g^2}{2}[-G(1, 3+4) - D_2(1)(\omega(3) + \omega(4) - 2\omega(3+4)) - D_2(1+2)(\omega(2) - \omega(1+2))]$$

as a sum of the double pomeron exchange coupled to $-(1/2)g^2 D_{20}(1)$ and a term $(1/2)g^2 D_2(1)$.

Evidently this result is trivially generalized for $f = D_2(i)$, $i = 2, 3, 4$ by simple permutation of indexes 1,2,3 and 4 and one may note that the separated terms found for the amplitude have the same momentum dependence of the corresponding zero order cancelled terms. Let us see if this is true also for the last three terms in (3.21).

Thus we consider the case $f = (1/2)g^2 D_2(1+2)$ for which we have

$$X = \frac{g^2}{2} D_{20} + \frac{g^2}{2} [g^2 N_c (V_0 - V_{12} - V_{34}) + 2\omega(1+2) - \sum \omega(i)] D_2(1+2) \quad (3.78)$$

In terms of W we have

$$V_0 D_2(1+2) = -W_2(1+2, 0, 3+4) \quad (3.79)$$

The bootstrap gives

$$g^2 N_c (V_{12} + V_{34}) D_2(1+2) = 2D_2(1+2)(2\omega(1+2) - \sum \omega(i)) \quad (3.80)$$

so that in terms of G we obtain

$$X = \frac{g^2}{2} [-D_{20}(1+2) + G(1+2, 3+4) - D(1+2)(4\omega(1+2) - \sum \omega(i))] \quad (3.81)$$

Again we see that the term with the double pomeron exchange coupled to $g^2(1/2)D_{20}(1+2)$ can be transformed into a triple pomeron vertex, essentially, into $-G(1+2, 3+4)$ term.

Finally we study a more complicated case with $f = (1/2)g^2 D_2(1+3)$. In this case we find

$$V_0 D_2(1+3) = -W_2(1+3, 0, 2+4) \quad (3.82)$$

Calculation of V_{12} or V_{34} applied to $D_2(1+3)$ is done using the formula (see appendix A.4)

$$V_{12} D_2(1+3) = W_2(2+4, 1, 3) + W_2(4, 2, 1+3) - W_2(3, 1+2, 4) - W_2(1+3, 0, 2+4) \quad (3.83)$$

and

$$V_{34} D_2(1+3) = W_2(2+4, 3, 1) + W_2(2, 4, 1+3) - W_2(1, 3+4, 2) - W_2(1+3, 0, 2+4) \quad (3.84)$$

Using these results one has

$$\begin{aligned} X = \frac{g^2}{2} & \left[D_{20}(1+3) + G(1, 2+4) + G(2, 1+3) + G(3, 2+4) + G(4, 1+3) \right. \\ & - G(1, 2) - G(3, 4) - G(1+3, 2+4) - D_2(1)(\omega(3+4) - \omega(3)) \\ & \left. - D_2(2)(\omega(3+4) - \omega(4)) - D_2(3)(\omega(1+2) - \omega(1)) - D_2(4)(\omega(1+2) - \omega(2)) \right] \end{aligned} \quad (3.85)$$

The result for $f = (1/2)g^2 D_2(1+4)$ is obtained from this after the permutation of 3 and 4.

Inspecting these results and comparing them with the form of our triple pomeron vertex, we see that only four terms

$$\frac{g^2}{2}[G(1,3) + G(1,4) + G(2,3) + G(2,4)] \quad (3.86)$$

are not changed under these transformations and thus correspond to a true triple pomeron interaction. All the rest can be transformed into terms which are essentially double pomeron exchange contribution. Conversely, one can eliminate terms from the double pomeron exchange substituting them by equivalent triple pomeron contributions.

The most radical result follows if one takes

$$f = D_{40}^{(0)}(D_{20} \rightarrow D_2) \quad (3.87)$$

In this case all the double exchange becomes cancelled and the whole amplitude is given by a sum of two terms (in an evident symbolic notation)

$$D_4^{(0)} = D_{40}^{(0)}(D_{20} \rightarrow D_2) + \int_0^Y G_2(Y-y)G_2(Y-y)\tilde{Z}D_2(y) \quad (3.88)$$

with a new vertex

$$\begin{aligned} \tilde{Z}D_2 = \frac{g^2}{2} & \left[G(1,3) + G(1,4) + G(2,3) + G(2,4) + G(1+2,3+4) \right. \\ & \left. - G(1,3+4) - G(2,3+4) - G(3,1+2) - G(4,1+2) \right] \end{aligned} \quad (3.89)$$

Comparing to (3.45) for the forward case, we observe that it coincides with the part $V(1234)$ of the vertex introduced in [20], being its leading part in the large N_c limit.

3.5 Relation to the dipole approach

According to the previous section we shall consider the DPE part completely transferred into the TPI one and study the triple pomeron vertex in the simpler form (3.45) for a non-forward direction. It is important to note that the coupling to BFKL pomerons of this vertex will lead to further simplifications. In fact in the coordinate space of 4 gluons, the dependence on only the sum of the momenta of two gluons, say, $1+2$, is translated into a factor $\delta^2(r_{12})$, so that the two gluons have to be taken at the same point. Since the wave functions $\Psi(r_1, r_2)$ and $\Psi(r_3, r_4)$ of the two final pomerons, coupled to the vertex, vanish if $r_1 = r_2$ and $r_3 = r_4$ respectively, all terms in eq.(3.45) which depend either only on the sum $1+2$ and/or only on the sum $3+4$ give zero, coupled to the two pomerons. Thus the effective triple pomeron vertex coupled to pomerons reads

$$\tilde{Z}D_2 = \frac{g^2}{2} \left[G(1,2+4,3) + G(1,2+3,4) + G(2,1+4,3) + G(2,1+3,4) \right] \quad (3.90)$$

Since both pomeron functions $\Psi(r_1, r_2)$ and $\Psi(r_3, r_4)$ are symmetric in their respective arguments, due to the positive signature of the pomeron, all terms in (3.90) give identical contributions and one has

$$\tilde{Z}D_2 = 2g^2G(1, 2 + 3, 4) \quad (3.91)$$

Moreover let us consider in the coordinate space the “improper part” of the function $G(r_1, r_2, r_3)$, which includes all the terms proportional to $\delta^2(r_{12})$ or/and $\delta^2(r_{23})$; since (3.91) is also proportional to $\delta^2(r_{23})$ we find that in the “improper part” at least three gluons, either 123 or 234, are to be taken at the same point in the transverse space. Then these terms will vanish due to the mentioned property of the pomeron wave function.

Thus only the contribution of the “proper part” is left and it is explicitly given by

$$G^{pr}(r_1, r_2, r_3) = -\frac{g^2 N_c}{8\pi^3} \frac{r_{13}^2}{r_{12}^2 r_{23}^2} D_2(r_1, r_3) \quad (3.92)$$

(with the regularization (3.51) implied).

Coupling this triple pomeron vertex to the two final pomerons, one obtains the following expression for the triple pomeron contribution to the diffractive (non-forward) amplitude:

$$D_4^{TPI}(Y) = -\frac{g^4 N_c}{4\pi^3} \int_0^Y dy \int d^2 r_1 d^2 r_2 d^2 r_3 \frac{r_{13}^2}{r_{12}^2 r_{23}^2} D_2(r_1, r_3; y) \Psi_1(r_1, r_2; Y - y) \Psi_2(r_2, r_3; Y - y) \quad (3.93)$$

If one writes the initial pomeron amplitude D_2 via the non-amputated function Ψ , which is the conformal invariant one, the amplitude reads

$$D_4^{TPI}(Y) = -\frac{g^4 N_c}{4\pi^3} \int_0^Y dy \int \frac{d^2 r_1 d^2 r_2 d^2 r_3}{r_{13}^2 r_{12}^2 r_{23}^2} \Psi_1(r_1, r_2; Y - y) \Psi_2(r_2, r_3; Y - y) r_{13}^4 \partial_1^2 \partial_3^2 \Psi(r_1, r_3; y) \quad (3.94)$$

In this form it is evident that the triple pomeron vertex is not symmetric with respect to the initial pomeron and two final ones: there appears an extra operator $r_{13}^4 \partial_1^2 \partial_3^2$ acting on the initial pomeron. Note that this operator is essentially a product of the Casimir operators of the conformal group for the holomorphic and antiholomorphic parts. Therefore it is convenient to expand the Ψ functions in terms of the conformal basis, introduced in the first chapter. Using (1.56) we do this for all the three pomerons in (3.94) and compute the action of the Casimir operators by means of (1.48). We then obtain, after the integration over y

$$D_4^{TPI}(Y) = -\frac{g^4 N_c}{4\pi^3} \sum_{\mu, \mu_1, \mu_2} \langle \mu | \Psi_0 \rangle \langle \mu_1 | \Psi_{10} \rangle \langle \mu_2 | \Psi_{20} \rangle \frac{e^{Y(\omega_{\mu_1} + \omega_{\mu_2})} - e^{Y\omega_\mu}}{\omega_{\mu_1} + \omega_{\mu_2} - \omega_\mu} \quad (3.95)$$

$$\frac{4\pi^8}{a_{n-1, \nu} a_{n+1, \nu}} \int \frac{d^2 r_1 d^2 r_2 d^2 r_3}{r_{13}^2 r_{12}^2 r_{23}^2} E_{\mu_1}(r_1, r_2) E_{\mu_2}(r_2, r_3) E_\mu(r_1, r_3)$$

In this form the triple pomeron contribution can be compared to the double dipole density found by Peschanski [31] in A.H.Mueller’s colour dipole approach. One observes that the two

expressions differ only in the sign and by a factor

$$\frac{4\pi^8}{a_{n-1,\nu}a_{n+1,\nu}}$$

which in our approach distinguishes the initial pomeron from the two final ones. The integral over the coordinates of the three pomerons is the same. So essentially the three-pomeron contribution to the diffractive amplitude found in our approach coincides with the double dipole density in the dipole approach.

However one should not forget that in our s -channel unitarity approach the TPI term (3.95) does not exhaust all the diffractive amplitude. In fact we have seen that

$$D_4^{(0)} = D_{40}(D_{20} \rightarrow D_2) + D_4^{TPI} \quad (3.96)$$

At high energies the TPI term behaves essentially as $s^{2\Delta}$ and the first one as s^Δ where Δ is the BFKL intercept. So one might think that the first term could be neglected. However the correct region of the validity of the leading log approximation, implicit in the hard pomeron theory, is $g^2 \ln s \sim 1$ when the two terms in (80) have the same order of magnitude. Moreover the BFKL separated term goes as $1/N_c$ and the TPI as $g^2 N_c$ and also these factors are of the same order. The dipole approach uses essentially the same leading log approximation but it leads to the double dipole density which coincides only with the TPI term in the s -channel unitarity approach and shows no trace of the first term. Therefore this fact points to certain differences between the two approaches.

3.6 Higher order densities in the dipole approach

Keeping in mind this substantial agreement between the results obtained in the s -channel unitarity approach and in the colour dipole picture for the two pomerons (dipoles) distribution, let us investigate the general structure of the higher order dipole densities given by the latter. To see if the previous equivalence found is maintained, one should study the multipomeron distribution in the former approach. In particular the first step would regard the study of the structure of the system of up to six reggeized gluons to look at the transition from 1 to 3 pomerons. In fact, as we shall show, this coupling does not directly appear in the standard dipole picture which instead leads to a set of fan diagrams with a single “elementary” three-pomeron coupling. If in such a context a future analysis will show the presence of pomeron vertices of order higher than 3 then an extension of the dipole picture, for example as the one proposed by Peshanski [31], should be required.

Following the approach of A.H. Mueller, as briefly reviewed in section 2.4, in the colour dipole formalism the k -fold inclusive dipole density is obtained as the k -th functional derivative of the functional $D\{u(r_i, r_f)\}$, taken at $u(r_i, r_f) = 1$ [11]. The arguments r_i and r_f are the dipole endpoints in the transverse plane. We remember here the simple equation the functional D satisfies

$$D(r_1, r_0, y, u) = u(r_1, r_0) e^{2y\omega(r_{10})}$$

$$+ \frac{g^2 N_c}{8\pi^3} \int_0^y dy' e^{2(y-y')\omega(r_{10})} \int d^2 r_2 \frac{r_{10}^2}{r_{12}^2 r_{20}^2} D(r_1, r_2, y', u) D(r_2, r_0, y', u) \quad (3.97)$$

Here r_1 and r_0 are the end points of the $q\bar{q}$ pair which determine the initial dipole; moreover we define $y = \ln z_{10}$ where z_{10} is the ratio of the scaling factors of the antiquark and of the quark; $\omega(r)$ is not a Fourier transform of the trajectory, but just $\omega(k)$ with k/m formally substituted by r/ϵ , where ϵ is an ultraviolet cutoff. This cutoff is also implied in the singular kernel of the integral operator in r_2 . Let us note that eq. (3.97) is compatible with the normalization condition $D(u=1) = 1$.

The k -th derivative of eq. (3.97) give an equation for the k -fold dipole density if computed for $u = 1$. For $k > 1$ we obtain

$$\begin{aligned} n_k(r_1, r_0; \rho_1, \dots, \rho_k; y) &= \frac{g^2 N_c}{8\pi^3} \int_0^y dy' e^{2(y-y')\omega(r_{10})} \int d^2 r_2 \frac{r_{10}^2}{r_{12}^2 r_{20}^2} n_k(r_1, r_2; \rho_1, \dots, \rho_k; y') + (r_1 \leftrightarrow r_0) \\ &+ \frac{g^2 N_c}{8\pi^3} \int_0^y dy' e^{2(y-y')\omega(r_{10})} \int d^2 r_2 \frac{r_{10}^2}{r_{12}^2 r_{20}^2} \sum_{l=1}^{k-1} (n_l(r_1, r_2; \rho_1, \dots, \rho_l; y') n_{k-l}(r_2, r_0; \rho_{l+1}, \dots, \rho_k; y') \\ &+ \text{symmetrization terms}) \end{aligned} \quad (3.98)$$

where the symmetrization terms (ST) are obtained from the explicitly shown one by taking all different divisions of arguments $\rho_1, \dots, \rho_k$ into two groups with l and $l - k$ arguments. In the $k = 1$ case an inhomogeneous term whose form is clear from (3.97) is present. One should note that the operator on the right-hand side acts nontrivially only on the first argument of the density n_k . Its action on the rapidity variable y , on the contrary, is rather simple and it is better treated in a transformed associated equation. Thus it is convenient to multiply the equation by $e^{-2y\omega(r_{10})}$, differentiate then with respect to y and pass to the j -space by the standard Mellin transformation. After having performed these operations one obtains

$$\begin{aligned} (j-1)n_k(r_1, r_0; \rho_1, \dots, \rho_k; j) &= \frac{g^2 N_c}{4\pi^3} \int d^2 r_2 L(r_1, r_2, r_{20}) n_k(r_1, r_2; \rho_1, \dots, \rho_k; y) \\ &+ \frac{g^2 N_c}{8\pi^3} \int \frac{dj_1 dj_2}{(2\pi i)^2 (j+1-j_1-j_2)} \int d^2 r_2 \frac{r_{10}^2}{r_{12}^2 r_{20}^2} \\ &\sum_{l=1}^{k-1} (n_l(r_1, r_2; \rho_1, \dots, \rho_l; j_1) n_{k-l}(r_2, r_0; \rho_{l+1}, \dots, \rho_k; j_2) + ST) \end{aligned} \quad (3.99)$$

where we introduced the BFKL kernel in the coordinate space

$$L(r_{12}, r_{20}) = \frac{r_{10}^2}{(r_{12}^2 + \epsilon^2)(r_{20}^2 + \epsilon^2)} - 2\pi \ln \frac{r_{10}}{\epsilon} [\delta^2(r_{12}) + \delta^2(r_{20})] \quad (3.100)$$

Comparing with (3.51) we see that it does not depend on ϵ and is ultraviolet stable. The solution of the equation (3.99) is quite easy to obtain if one describes the dependence of the densities on

their first two arguments in terms of the conformal basis:

$$n_k(r_1, r_0) = \sum_{\mu} E_{\mu}(r_1, r_0) n_k^{\mu} \quad (3.101)$$

Here we have suppressed all other arguments in n_k irrelevant for the time being. The densities n_k^{μ} in a given conformal state are obtained from $n_k(r_1, r_0)$ by the inverse transformation which follows from orthogonality property (1.52) and a relation between $E_{n,\nu}$ and $E_{-n,-\nu}$ (see [27])

$$n_k^{\mu} = \int \frac{d^2 r_1 d^2 r_0}{r_{10}^4} E_{\mu}^*(r_1, r_0) n_k(r_1, r_0) \quad (3.102)$$

Therefore, to pass to the conformal basis, we integrate eq. (3.99) over r_1 and r_0 as indicated in (3.102). The first term on the right-hand side can be simplified due to the property of the BFKL kernel

$$\frac{g^2 N_c}{4\pi^3} \int d^2 r_2 L(r_{12}, r_{20}) E_{\mu}(r_1, r_2) = \omega_{\mu} E_{\mu}(r_1, r_0) \quad (3.103)$$

Therefore after the integration we obtain

$$(j-1-\omega_{\mu}) n_k^{\mu}(\rho_1, \dots, \rho_k, j) = \frac{g^2 N_c}{8\pi^3} \int \frac{dj_1 dj_2}{(2\pi i)^2 (j+1-j_1-j_2)} \int \frac{d^2 r_1 d^2 r_2 d^2 r_0}{r_{12}^2 r_{20}^2 r_{10}^2} E_{\mu}^*(r_{10}) \sum_{l=1}^{k-1} (n_l(r_1, r_2; \rho_1, \dots, \rho_l; j_1) n_{k-l}(r_2, r_0; \rho_{l+1}, \dots, \rho_k; j_2) + ST) \quad (3.104)$$

To find the final form of the equation we have only to present also the densities n_l and n_{k-l} as functions of their first arguments in the form (3.102). Then one has

$$(j-1-\omega_{\mu}) n_k^{\mu}(\rho_1, \dots, \rho_k, j) = \int \frac{dj_1 dj_2}{(2\pi i)^2 (j+1-j_1-j_2)} \sum_{\mu_1, \mu_2} V_{\mu, \mu_1 \mu_2} \sum_{l=1}^{k-1} (n_l^{\mu_1}(\rho_1, \dots, \rho_l; j_1) n_{k-l}^{\mu_2}(\rho_{l+1}, \dots, \rho_k; j_2) + ST) \quad (3.105)$$

where

$$V_{\mu \mu_1 \mu_2} = \frac{g^2 N_c}{8\pi^3} \int \frac{d^2 r_1 d^2 r_2 d^2 r_0}{r_{12}^2 r_{20}^2 r_{10}^2} E_{\mu}^*(r_{10}) E_{\mu_1}(r_{12}) E_{\mu_2}(r_{20}) \quad (3.106)$$

is just one half of the three-pomeron vertex introduced by Peschanski. The general structure of this equation allows one immediately to imagine a tree-like expansion of the colour dipole densities as we shall show more in detail. At each tree node a vertex (3.106) will be present. The integral of its explicit expression has been calculated [28, 58] with the leading quantum numbers $n = \nu = 0$ for the related three pomerons and it turns out to have a quite huge value (7766.679). Of course there will be present some correcting factors but still this means that such a perturbative expansion in pomeron vertices may need a complete resummation.

Eq. (3.105) allows to obtain successively dipole densities for any number of dipoles starting

from the lowest order one-dipole density, for which

$$n_1^\mu(\rho) = \frac{E_\mu^*(\rho)}{j-1-\omega_\mu} \frac{1}{\rho^4} \quad (3.107)$$

(we recall that in this notation ρ includes two endpoints of the dipole ρ_i and ρ_f ; $\rho^2 \equiv \rho_{if}^2$).

Putting this into (3.105) for $k=2$ and integrating over j_1 and j_2 we arrive at the expression obtained by Peschanski

$$n_2^\mu(\rho_1, \rho_2; j) = \frac{1}{\omega - \omega_\mu} \sum_{\mu_1, \mu_2} \frac{1}{\omega - \omega_{\mu_1} - \omega_{\mu_2}} V_{\mu, \mu_1, \mu_2} E_{\mu_1}^*(\rho_1) E_{\mu_2}^*(\rho_2) \frac{1}{\rho_1^4 \rho_2^4} \quad (3.108)$$

where $\omega = j-1$. (To compare with [31] one should take into account that factors $1/(2a_\mu)$ are included in the definition of sums over μ 's in our notation).

As a further step we study the density for three dipoles. Eq. (3.105) for $k=3$ reads

$$n_3^\mu(\rho_1, \rho_2, \rho_3; j) = \frac{1}{\omega - \omega_\mu} \int \frac{dj_1 dj_2}{(2\pi i)^2 (j+1-j_1-j_2)} \sum_{\mu_1, \mu_2} V_{\mu, \mu_1 \mu_2} (n_1^{\mu_1}(\rho_1; j_1) n_2^{\mu_2}(\rho_2, \rho_3; j_2) + ST) \quad (3.109)$$

It is instructive to study explicitly this term inserting the expressions for $n_1^{\mu_1}$ and $n_2^{\mu_2}$ obtained earlier. Then we get, after integrations over j_1 and j_2 :

$$n_3^\mu(\rho_1, \rho_2, \rho_3; j) = \frac{1}{\omega - \omega_\mu} \sum_{\mu_1, \mu_2, \mu_3, \mu_4} V_{\mu, \mu_1 \mu_2} V_{\mu_2 \mu_3 \mu_4} \frac{1}{(\omega - \omega_{\mu_1} - \omega_{\mu_2})(\omega - \omega_{\mu_1} - \omega_{\mu_3} - \omega_{\mu_4})} E_{\mu_1}^*(\rho_1) E_{\mu_3}^*(\rho_2) E_{\mu_4}^*(\rho_3) \frac{1}{\rho_1^4 \rho_2^4 \rho_3^4} \quad (3.110)$$

To this term we have to add terms which symmetrize in the three dipoles.

Studying (3.110) we see that it corresponds to the picture when first the initial pomeron splits into two pomerons, 1 and 2, and afterwards the pomeron 2 splits into pomerons 3 and 4. One does not find here a local vertex for the transition of the initial pomeron into three final ones. It is not difficult to see under which condition one would get such a local vertex. If we forget about the dependence of the second denominator on μ_2 , then one can sum over μ_2 . Using the completeness relation (1.50) one obtains

$$\sum_{\mu_2} V_{\mu \mu_1 \mu_2} V_{\mu_2 \mu_3 \mu_4} = \left(\frac{g^2 N_c}{8\pi^3} \right)^2 \int \frac{d^2 r_1 d^2 r_2 d^2 r_3 d^2 r_0}{r_{12}^2 r_{10}^2 r_{23}^2 r_{30}^2} E_\mu^*(r_1, r_0) E_{\mu_1}(r_1, r_2) E_{\mu_3}(r_2, r_3) E_{\mu_4}(r_3, r_0) \quad (3.111)$$

which is just the vertex from one to three pomerons introduced by Peschanski. However, the described summation is not possible due to the second denominator. It implies that the pomeron 2 has to evolve in y from the point of its formation from the initial pomeron up to the point of its splitting into the final pomerons 3 and 4.

Thus our conclusion is that the vertex for transition from 1 to k pomerons introduced by

Peschanski, in fact, does not appear in the solution of the Mueller equation (3.98) for the k -fold density, which rather corresponds to a set of all fan diagrams with only the triple pomeron coupling. Absence of higher-order couplings can be directly traced to the structure of the equation (3.97) for the generating functional D , quadratic in D , which in fact means that the basic vertex changes one dipole into two.

The general structure of higher order vertices introduced by Peschanski hence refers to an extension, which he proposes, to the dipole model. In his work the possibility of a splitting of one dipole into many dipoles is considered, going therefore beyond the standard dipole model. His idea is related to the fact that in general a contribution to the transition from 1 to p pomeron given by the fan diagrams would require a large rapidity interval at each splitting step in order to ensure the validity of the Regge approximation. The derived effective field theory is therefore more complex but it should be characterized by a strong symmetry since the vertices are related each other by integral relations, as for example the one in eq. (3.111).

A general analysis of the n -reggeon amplitudes approach, based on s -channel unitarity, is not known yet, even if some work in this direction is in progress. Any advance in this direction will be of great interest to understand the Regge limit effective behaviour of QCD.

To conclude this section we shall try to analyze in more details the picture one derives from the triple pomeron interaction structure so far verified. It has been shown that at asymptotic energies the higher-order densities in the dipole approach correspond to the standard Regge-Gribov picture, in the tree approximation (fan diagrams), with only the triple pomeron interaction, which however has a highly complicated non-local form. Indeed, the triple pomeron interaction present in (3.110) corresponds to a structure

$$T = \frac{g^2 N_c}{8\pi^3} \int \frac{d^2 r_1 d^2 r_2 d^2 r_3}{r_{12}^2 r_{23}^2 r_{31}^2} \tilde{G}_3(r_1, r_2; r'_1, r'_2) \tilde{G}_1(r_2, r_3; r'_2, r'_3) \tilde{G}_2(r_3, r_1; r'_3, r'_1) \quad (3.112)$$

where $\tilde{G}_i$, $i = 1, 2, 3$ are Green functions of the interacting pomerons defined as

$$\tilde{G}(r_1, r_2; r'_1, r'_2) = \sum_{\mu} \frac{E_{\mu}(r_1, r_2) E_{\mu}^*(r'_1, r'_2)}{\omega - \omega_{\mu}} \quad (3.113)$$

They are not the physical BFKL Green functions. The latter include an extra factor depending on μ :

$$G(r_1, r_2; r'_1, r'_2) = \frac{1}{4\pi^8} \sum_{\mu} a_{n+1, \nu} a_{n-1, \nu} \frac{E_{\mu}(r_1, r_2) E_{\mu}^*(r'_1, r'_2)}{\omega - \omega_{\mu}} \quad (3.114)$$

However in the limit $s \rightarrow \infty$ only the lowest conformal weights contribute $n = \nu = 0$ for which $a_{n \pm 1, \nu} = 2\pi^4$ and (3.113) and (3.114) coincide. Then we can forget about tildes in (3.112).

We transform the Green functions to given total momenta of the pomerons presenting

$$G_3(r_1, r_2; r'_1, r'_2) = \int \frac{d^2 l_3}{(2\pi)^2} e^{il_3(R_3 - R'_3)} G_{l_3}(r_{12}, r'_{12}) \quad (3.115)$$

where $R_3 = (1/2)(r_1 + r_2)$ and similarly for the two other Green functions. Introducing $R =$

$r_1 + r_2 + r_3$ we transform the integration over the coordinates as follows

$$\int d^2r_1 d^2r_2 d^2r_3 = \int d^2R d^2r_{12} d^2r_{23} d^2r_{31} \delta^2(r_{12} + r_{23} + r_{31}) \quad (3.116)$$

The coordinates themselves are $r_1 = (1/3)(R - r_{21} - r_{31})$ etc., where from we find

$$R_1 = (1/6)(2R - r_{12} - r_{13}), \quad R_2 = (1/6)(2R - r_{21} - r_{23}), \quad R_3 = (1/6)(2R - r_{31} - r_{32}) \quad (3.117)$$

and

$$i \sum_{j=1}^3 l_j R_j = i(1/3)R \sum_{j=1}^3 l_j - i(1/6)(r_{12}l_{12} + r_{23}l_{23} + r_{31}l_{31}) \quad (3.118)$$

where we denoted $l_{12} = l_1 - l_2$ etc. The integral over R gives $9(2\pi)^2 \delta^2(l_1 + l_2 + l_3)$. Presenting the remaining δ -function in (3.116) as an integral over an auxiliary momentum q we find an expression (for fixed l_1, l_2 and l_3)

$$\begin{aligned} & \frac{g^2 N_c}{8\pi^3} \int \frac{d^2q}{(2\pi)^2} \frac{d^2r_{12} d^2r_{23} d^2r_{31}}{r_{12}^2 r_{23}^2 r_{31}^2} \exp \left(ir_{12} \left(q - \frac{1}{6} l_{12} \right) + ir_{23} \left(q - \frac{1}{6} l_{23} \right) + ir_{31} \left(q - \frac{1}{6} l_{31} \right) \right) \\ & G_{l_3}(r_{12}, r'_{12}) G_{l_1}(r_{23}, r'_{23}) G_{l_2}(r_{31}, r'_{31}) \end{aligned} \quad (3.119)$$

At this point we recall the expression for the BFKL Green function with a fixed total momentum:

$$G_l(r, r') = \frac{1}{(2\pi)^4} \int \frac{\nu^2 d\nu}{(\nu^2 + 1/4)^2} s^{\omega(\nu)} E_\nu^{(l)}(r) E_\nu^{(l)}(r') \quad (3.120)$$

where

$$E_\nu^{(l)}(r) = \int d^2R \exp(ilR) \left(\frac{r}{|R + r/2| |R - r/2|} \right)^{1+2i\nu} \quad (3.121)$$

and where we retained only the dominant isotropic term. At $s \rightarrow \infty$ the vicinity of $\nu = 0$ gives the dominant contribution. If $l \neq 0$ then the integral in (3.121) converges at large R and we can take the functions E out of the integral over ν at $\nu = 0$. Taking then the asymptotics of the remaining integral, we find

$$G_l(r, r') \simeq \frac{1}{2\pi^4} s^\Delta \frac{\sqrt{\pi}}{(a \ln s)^{3/2}} E_0^{(l)}(r) E_0^{(l)}(r') \quad (3.122)$$

where $\Delta = \omega_{n=0, \nu=0}$ is the BFKL intercept and $a = 7g^2 N_c \zeta(3)/(2\pi^2)$. As we see, the Green function asymptotically factorizes in r and r' . This means that we obtain a quantum field theory of pomerons with a propagator

$$P(y, l) = \frac{2}{\pi^2} e^{y\Delta} \frac{\sqrt{\pi}}{(ay)^{3/2}} \quad (3.123)$$

(not really depending on the momentum l) and an interaction vertex

$$V(l_1, l_2, l_3) = \frac{9g^2 N_c}{8\pi^3} \int \frac{d^2q}{(2\pi)^8} \frac{d^2r_{12} d^2r_{23} d^2r_{31}}{r_{12}^2 r_{23}^2 r_{31}^2}$$

$$\exp\left(ir_{12}\left(q - \frac{1}{6}l_{12}\right) + ir_{23}\left(q - \frac{1}{6}l_{23}\right) + ir_{31}\left(q - \frac{1}{6}l_{31}\right)\right) E_0^{(l_3)}(r_{12})E_0^{(l_1)}(r_{23})E_0^{(l_2)}(r_{31}) \quad (3.124)$$

The vertex factorizes under the sign of the integration over q :

$$V(l_1, l_2, l_3) = \frac{9g^2 N_c}{8\pi^3} \int \frac{d^2 q}{(2\pi)^2} J(l_3, q - \frac{1}{6}l_{12}) J(l_1, q - \frac{1}{6}l_{23}) J(l_2, q - \frac{1}{6}l_{31}) \quad (3.125)$$

where

$$J(l, q) = \int \frac{d^2 r}{(2\pi)^2 r^2} e^{iqr} E_0^{(l)}(r) = \int \frac{d^2 p}{2\pi p} \frac{1}{|p - q + l/2||p - q - l/2|} \quad (3.126)$$

Note that for $l = 0$ this derivation is incorrect. Calculations show that in this case eq. (3.125) for the vertex remains valid with

$$J(0, q) = \frac{1}{9q} \quad (3.127)$$

However the Green function (3.120) at $l = 0$ has an asymptotics

$$G_0(r, r') \simeq \frac{1}{2\pi^2} s^\Delta \sqrt{\frac{\pi}{a \ln s}} r r' \exp\left(-\frac{\ln^2(r/r')}{a \ln s}\right) \quad (3.128)$$

which is evidently no more factorizable in r and r' .

Chapter 4

Conclusions

As stated in the introduction, the thesis has been devoted to the study of some issues relevant for the hard pomeron model. Both theoretical and phenomenological aspects have been considered.

From a phenomenological point of view in the second chapter an extension to include the running coupling for the hard pomeron is studied. The leading singularity in this way is softened and once the infrared sector is safely included in the model one can avoid also the problems due to the diffusion mechanism which spoil out the validity of the pure BFKL pomeron. The model, which preserves the gluon reggeization property by means of the bootstrap condition and the right gluon distribution behaviour in the DLLA, has been studied both numerically and in some aspects analytically. Numerical calculations were addressed to the computation of the spectrum and eigenfunctions of the Schrödinger-like equation for the pomeron. The spectrum has shown a discrete part and for the wide parameter range two negative “energies” states corresponding to two supercritical pomeron states. The slope of each singularity has been calculated with a perturbative approach. The sensitivity of the model is mainly with respect to only one infrared parameter, a mass which we chose to fix forcing a value for the slope of the leading singularity. We have applied such a model to the process of two virtual photon scattering and to see some unitarization effects we have considered in the large N_c limit a multi-scattering (eikonal) formulation which have shown small unitarity corrections of such a kind to the cross sections. Triple pomeron interactions, required by unitarity, are not taken into account in these calculations and larger unitarity effects are therefore not excluded. Similar considerations were applied to a toy model, with a Gaussian colour dipole distribution, for the proton.

An application of this model to the inclusive jet production has been considered. A correct asymptotic behaviour of the differential cross section for large jet transverse momenta has been found. In this case one must note that in the BFKL approach already some unitarity correction, which have lead to modify the BFKL equation with a non-linear contribution generating fan diagrams, as well as coherence effects were, for example, invoked. Since the leading singularity is a pole, the y dependence in the cross section depends on the inclusion of the non-leading states of the spectrum. Here we calculated the contribution due to the two supercritical pomerons and for a better estimate a much larger part of the spectrum depending on the energy, should be considered.

The contribution from the whole spectrum have been included in the analysis of the evolu-

tion of the gluon density and of the proton singlet structure function. The initial proton gluon distribution in rapidity has been found by matching the related singlet structure function. The main problem has been to find a way to couple the gluon to the photon taking in account the introduced running of the coupling. A phenomenological approach has been chosen, based on the requirement that for the structure function in the DLLA the analytical asymptotic behaviour should match the one calculated using the DGLAP quark distributions. The numerical calculations have still shown a steep increase with decreasing x of the gluon distribution, but with a better behaviour than for a BFKL pomeron.

In the theoretical part of the thesis we deal with the unitarity problem. At the basis of this contribution are the works of J.Bartels et al. where a program to ensure unitarity requirements has been developed and the analysis of the lower order corrections, up to 4 gluons in the t -channel, started. The idea of studying these corrections in the large N_c limit, a program initiated by M.Braun, is pursued in order to avoid the otherwise complicated colour structure. The nice property of this limit is that one does not lose many important features of the underlying interactions and it is also possible to directly compare the obtained results with some recent developments derived in the colour dipole picture. In particular the system of four gluons has been studied for a non-zero momentum flow in the t channel in the leading and next-to-leading order in $1/N_c$. The solution has been shown to have in general double pomeron exchange and triple pomeron interaction contributions. The latter is written in terms of a three point function which is by itself infrared stable and conformal invariant. Following the idea of Bartels and Wuesthoff it is shown the possibility of eliminating the DPE terms in the solution so that the unitarity correction to this order may depend only on a particular three pomeron vertex, plus some single pomeron terms. There is complete agreement with the Bartels et al. works once the large N_c limit is taken into account. After coupling this vertex to pomerons one can easily compare the obtained two pomeron amplitude with the double dipole colour density analyzed by R.Peschanski in the standard dipole picture. One finds a substantial agreement for the triple pomeron interaction form, apart from an asymmetry factor and the previous mentioned single pomeron terms. The properties of higher order dipole densities, as defined in the A.H. Mueller formulation of the colour dipole model, are subsequently studied. A picture is found where in the rapidity evolution a pomeron may split into two pomerons, thus leading to a fan diagrams description with only three pomeron vertices involved. The resulting pomeron effective field theory would hence have a strong signature. Therefore a corresponding analysis is needed in the reggeon approach based on s -channel unitarity. The analysis starting from a system of 5 and 6 reggeized gluons and in general of higher order distributions will show if this picture is correct or one has to deal with higher order effective pomeron vertices. In such a case one should also consider extension in the colour dipole picture, as for example proposed by R.Peschanski. In any case the structure of a possible effective theory of QCD in the Regge limit will then be strongly restricted.

Acknowledgements

I would like to express my deep gratitude to my supervisor Professor Giovanni Venturi for his support, interest and encouragement and for countless stimulating discussions in different fields of Physics.

I am sincerely and infinitely grateful also to Professor Mikhail Braun, for his guidance and help in the last years. His ideas have had a strong influence and most of the work present in the thesis is inspired and arises from his collaboration.

It was a great experience to visit the University of Santiago de Compostela and I would like to thank Profs. C. Pajares and L. Miramontes for their hospitality.

Furthermore I thank Prof. R. Peschanski for pleasant and useful discussions.

Finally a special thank to Fabio Finelli for many discussions and a fruitful collaboration.

Appendix A

A.1 Asymptotic behaviour of the wave functions.

A.1.1 The momentum space behaviour

Let us consider the asymptotic expression for the gluon Regge trajectory given in (2.11)

$$\omega(q) \underset{q^2 \gg m^2}{\sim} -\frac{N_c}{2b} \ln \frac{\ln q^2}{\ln m^2} \quad (\text{A.1})$$

Hence, for $N_c = 3$, one can write the asymptotic form of the pomeron non-homogeneous equation at high momentum for the semi-amputated wave function ψ

$$\ln \frac{\ln q^2}{\ln m^2} \psi(q) + \frac{am^2 \ln m^2}{q^2 \ln q^2} = \tilde{\epsilon} \psi(q) + \frac{1}{\pi} \int \frac{d^2 q' \psi(q')}{[(q - q')^2 + m^2] \ln[(q - q')^2 + m^2]} \quad (\text{A.2})$$

where we have put $E = (3/b)\tilde{\epsilon}$ (we consider the case $m = m_1$ for simplicity) Let us study the last (integral) term, which we denote by D . Changing the variable $q' = |q|\kappa$ we present it in the form

$$\begin{aligned} D &= \frac{1}{\pi} \int \frac{d^2 \kappa \psi(|q|\kappa)}{[(n - \kappa)^2 + m^2/q^2] \ln[q^2(n - \kappa)^2 + m^2]} \\ &= \frac{1}{\pi} \int \frac{d^2 \kappa \psi(|q|(n + \kappa))}{(\kappa^2 + m^2/q^2) \ln(q^2 \kappa^2 + m^2)} \end{aligned} \quad (\text{A.3})$$

with $n^2 = 1$. In the last form it is evident that the leading terms at $q \rightarrow \infty$ come from the integration region of small κ . Then we split the total κ space into two parts: $\kappa > \kappa_0$ and $\kappa < \kappa_0$ where κ_0 is a small number

$$\kappa_0 \ll 1 \quad (\text{A.4})$$

The contributions from these two parts we denote as D_1 and D_2 , respectively. Our first task is to show that D_2 cancels the kinetic term in (A.2) irrespective of the asymptotics of $\psi(q)$.

With small enough κ_0 and for a "good enough" ψ

$$\psi(|q|(n + \kappa)) \simeq \psi(|q|n) = \psi(q) \quad (\text{A.5})$$

so that we can take it out of the integral over κ in D_2 . Actually (A.5) is our definition of a "good" function, so that we shall have to check if this condition is indeed satisfied for the found

asymptotic $\psi(q)$.

With (A.5) D_2 simplifies to

$$D_2 = \frac{\psi(q)}{\pi} \int_0^{\kappa_0^2 q^2} \frac{d^2 q'}{(q'^2 + m^2) \ln(q'^2 + m^2)} = \psi(q) \ln \frac{\ln(\kappa_0^2 q^2 + m^2)}{\ln m^2} \quad (\text{A.6})$$

As $q \rightarrow \infty$ we assume that also

$$\kappa_0 q \rightarrow \infty \quad (\text{A.7})$$

Evidently this condition fixes the manner in which κ_0 goes to zero as q turns large. This should be taken into account when verifying condition (A.4). Then (A.6) gives

$$D_2 = \psi(q) \ln \frac{\ln \kappa_0^2 q^2}{\ln m^2} \quad (\text{A.8})$$

We have also

$$\ln \ln \kappa_0^2 q^2 = \ln(\ln q^2 + \ln \kappa_0^2)$$

But according to (A.7)

$$\ln q^2 \gg |\ln \kappa_0^2|$$

so that we have

$$\ln \ln \kappa_0^2 q^2 = \ln \ln q^2 + \frac{\ln \kappa_0^2}{\ln q^2} + \dots$$

Then our final result is

$$D_2 = \psi(q) \left(\ln \frac{\ln q^2}{\ln m^2} + O(1/\ln q^2) \right) \quad (\text{A.9})$$

The first term exactly cancels the kinetic energy (first) term in the pomeron equation. The correction term in (A.9) evidently is much smaller than $\tilde{\epsilon}\psi(q)$, since the factor which multiplies the function ψ goes to zero in (A.9). So we can safely neglect it.

As a result, the asymptotic equation becomes

$$\frac{am^2 \ln m^2}{q^2 \ln q^2} = \tilde{\epsilon}\psi(q) + D_1 \quad (\text{A.10})$$

with the term D_1 given by

$$D_1 = \frac{1}{\pi} \int \frac{d^2 q' \psi(q') \theta((q - q')^2 - \kappa_0^2 q^2)}{(q - q')^2 \ln(q - q')^2} \quad (\text{A.11})$$

where we have omitted the m^2 terms in the denominator because of (A.7).

We consider the function $\chi(q)$ defined by

$$\chi(q) = \psi(q) q^2 \ln q^2 \quad (\text{A.12})$$

Multiplying (A.10) by $q^2 \ln q^2$ we find an equation for χ

$$am^2 \ln m^2 = \tilde{\epsilon} \chi(q) + \frac{1}{\pi} \int \frac{d^2 q' \chi(q') \theta((q - q')^2 - \kappa_0^2 q^2)}{q'^2 \ln q'^2} \frac{q^2 \ln q^2}{(q - q')^2 \ln(q - q')^2} \quad (\text{A.13})$$

At this point we make a second assumption. Namely we assume that in the integral term of (A.13) values $m \ll q' \ll q$ give the dominant contribution (as typical for logarithmic integrals). Of course, this assumption is also to be checked for the final asymptotics. With this assumption, we can forget about the θ function and also put the last factor equal to unity in the integral term of (A.13). We obtain

$$am^2 \ln m^2 = \tilde{\epsilon} \chi(q) + \int_0^{q^2} \frac{dq'^2 \chi(q')}{q'^2 \ln q'^2} \quad (\text{A.14})$$

Differentiating with respect to q^2

$$\tilde{\epsilon} \frac{d\chi}{dq^2} = -\frac{\chi}{q^2 \ln q^2} \quad (\text{A.15})$$

or

$$\tilde{\epsilon} \frac{d\chi}{d \ln \ln q^2} = -\chi \quad (\text{A.16})$$

with a solution

$$\chi(q) = A \exp \left(-\frac{\ln \ln q^2}{\tilde{\epsilon}} \right) = A (\ln q^2)^{-1/\tilde{\epsilon}} \quad (\text{A.17})$$

The initial function $\psi(q)$ has the asymptotics

$$\psi(q) = \frac{A}{q^2} (\ln q^2)^\beta \quad (\text{A.18})$$

where

$$\beta = -1 - \frac{1}{\tilde{\epsilon}} = -1 - \frac{3}{bE} \quad (\text{A.19})$$

Now we must check that our assumptions are indeed fulfilled for the found asymptotics.

Let us begin with the second assumption that the values $m \ll q' \ll q$ give the bulk of the contribution to the integral in (A.13). Evidently, in order that the integral be dominated by large values of q' , it should diverge as $q \rightarrow \infty$. This leads to the condition

$$E < 0, \quad \beta > -1 \quad (\text{A.20})$$

So our asymptotics can only be valid for negative energies, that is, for bound states.

Now for the second part of this assumption. To prove that values $q' \ll q$ dominate we shall calculate the contribution from the region $q' \gg q$ and show that it is smaller. The corresponding integral is

$$I = A q^2 \ln q^2 \int_{q^2}^{\infty} dq'^2 (\ln q'^2)^{\beta-1} / q'^4 \quad (\text{A.21})$$

In terms of $x = \ln q'^2$

$$I = q^2 \ln q^2 \int_{\ln q^2}^{\infty} dx x^{\beta-1} \exp(-x) \quad (\text{A.22})$$

The integral over x can be developed in an asymptotic series in $1/\ln q^2$:

$$\int_{\ln q^2}^{\infty} dx x^{\beta-1} \exp(-x) = - \int_{\ln q^2}^{\infty} x^{\beta-1} d \exp(-x) = \frac{(\ln q^2)^{\beta-1}}{q^2} + (\beta-1) \int_{\ln q^2}^{\infty} dx x^{\beta-2} \exp(-x) = \dots \quad (\text{A.23})$$

From this we conclude that the integral I has the asymptotics

$$I = A(\ln q^2)^{\beta} \quad (\text{A.24})$$

to be compared to the contribution from the region $q' \ll q$ which behaves as $(\ln q^2)^{\beta+1}$. We see that we have lost one power of $\ln q^2$, so that the region $q' \gg q$ indeed can be neglected.

Now to the assumption (A.5). We have explicitly

$$\begin{aligned} \psi(|q|(n+\kappa)) &= (A/q^2)(n+\kappa)^{-2}(\ln q^2(n+\kappa)^2)^{\beta} = \\ &= (A/q^2)(1-2n\kappa-\kappa^2+4(n\kappa)^2)(\ln q^2)^{\beta}(1+(\beta(2n\kappa+\kappa^2)/\ln q^2)) \end{aligned} \quad (\text{A.25})$$

and it is evident that (A.11) is satisfied with the behaviour of κ as indicated in (A.7). Indeed take $\kappa \sim q^{-\delta}$ with $\delta < 1$. Then $q\kappa \sim q^{1-\delta} \rightarrow \infty$ and all the correcting terms in (A.25) have the order $q^{-\delta}$.

Note that this does not mean that (A.5) is quite obvious. It is not valid for, say, the exponential function. In fact we have

$$\exp(aq^2(n+\kappa)^2) = \exp(aq^2) \exp(aq^2(2n\kappa+\kappa^2))$$

and since $q\kappa$ is large the second factor cannot be neglected.

In conclusion, we have verified that our assumptions are fulfilled and therefore the asymptotics (A.18) is correct for negative “energies”.

A.1.2 The coordinate space behaviour

We are also interested in the coordinate space behaviour of the semi-amputated wave function

$$\psi(r) = \int_0^{\infty} \frac{q dq}{2\pi} J_0(qr) \psi(q) \quad (\text{A.26})$$

To estimate $\psi(r)$ for $r \rightarrow 0$ we make use of the asymptotic momentum behaviour found previously (A.18), so we can write

$$\psi(r) \approx \int_0^{q_0} \frac{q dq}{2\pi} J_0(qr) \psi(q) + \int_{q_0}^{\infty} \frac{q dq}{2\pi} J_0(qr) \frac{\ln(q^2)^{\beta}}{q^2} \quad (\text{A.27})$$

The first integral for $r \rightarrow 0$ is finite; on the other hand the second integral results to be not bounded. Infact using the integration variable $y = qr$ we split the y integration region in two

parts, $q_0 r < y < y_0 \ll 1$ and $y \geq y_0$; defining the two contribution I_1 and I_2 respectively, we have

$$I_1 \sim \int_{q_0 r}^{y_0} \frac{dy}{y} \left(\ln \frac{y}{r} \right)^\beta \underset{r \rightarrow 0}{\sim} \left(\ln \frac{1}{r} \right)^{\beta+1} \quad (\text{A.28})$$

and

$$I_2 \sim \int_{y_0}^{\infty} \frac{dy}{y} J_0(y) \left(\ln y + \ln \frac{1}{r} \right)^\beta \underset{r \rightarrow 0}{\sim} \left(\ln \frac{1}{r} \right)^\beta \quad (\text{A.29})$$

So the asymptotic small r behaviour will be divergent, in particular

$$\psi(r) \underset{r \rightarrow 0}{\sim} \left(\ln \frac{1}{r} \right)^{\beta+1} \quad (\text{A.30})$$

A.2 Two gluons coupling to quark loop for non-zero transfer momentum

Consider a $q\bar{q}$ loop for the scattering of a virtual photon $\gamma^*(q) + \dots \rightarrow \gamma^*(q+l) + \dots$, $q^2 = -Q^2$. The momentum transfer l is taken to be purely transversal. We are interested in a multi-discontinuity amplitude so that at the lowest order one retains at each discontinuity a simple $q\bar{q}$ state with the quark and antiquark on shell (due to unitarity) and then with their propagators substituted by the corresponding delta distributions. The calculations are quite straightforward.

For example let us consider the most involved case of a transverse polarized photon and the gluons (with colours a_1 and a_2) coupled one (with momentum q_1) to the quark (with momentum k_1) and the other to the anti-quark (momenta q_2 and k_2 respectively) so that $q = k_1 + k_2$. In this case the propagators for the quark with momentum $k_1 + q_1$ and the anti-quark with momentum k_2 have to be replaced by the mass-shell conditions.

In the high energy limit only the longitudinal component of the gluons polarization give a contribution. It is convenient to work in the light cone coordinates and therefore one has for the trace present in the amplitude the following expression

$$N_{\perp} = \frac{\delta_{a_1 a_2}}{2} g_{\alpha\beta} \text{Tr} \left[\gamma_{\perp}^{\alpha} (m_f + \gamma \cdot k_1) \gamma_{-} (m_f + \gamma \cdot (k_1 + q_1)) \gamma_{\perp}^{\beta} (m_f - \gamma \cdot (k_2 + q_2)) \gamma_{-} (m_f - \gamma \cdot k_2) \right] \quad (\text{A.31})$$

After some algebra and performing the trace one obtains

$$N_{\perp} = -8\delta_{a_1 a_2} q_{-}^2 \left\{ m_f^2 + [\alpha^2 + (1 - \alpha)^2] k_{1\perp} \cdot (k_1 + q_1)_{\perp} - \alpha^2 k_{1\perp} \cdot l_{\perp} \right\} \quad (\text{A.32})$$

where α is defined such that $k_1 = \alpha q_{-}$. The propagators contain in the denominator two terms which can be written, defining $Q^2 = -q^2$, as

$$\begin{aligned} D_1 = (k_1^2 - m_f^2) &= -\frac{1}{1 - \alpha} \left[Q^2 \alpha (1 - \alpha) + (m_f^2 + k_{1\perp}^2) \right] \\ D_2 = [(k_2 + q_2)^2 - m_f^2] &= -\frac{1}{\alpha} \left\{ \alpha (1 - \alpha) (Q^2 + l^2) + m_f^2 + [(k_1 + q_1)_{\perp} - \alpha l_{\perp}]^2 \right\} \end{aligned} \quad (\text{A.33})$$

Taking also into account the contribution of the δ distributions, $1/4q_{-}\alpha(1-\alpha)$, one easily obtains the full expression. Similar expressions can be obtained for different photon polarizations and configurations of gluons.

Thus the calculations give for the function $f(1, 2)$ corresponding to the loop with gluon 1 attached to q and gluon 2 attached to $\bar{q}$ the following expression

$$f(k_1, k_2) = e_f^2 \int_0^1 d\alpha \int \frac{d^2 k}{(2\pi)^3} \frac{N}{D} \quad (\text{A.34})$$

where now we use $\mathbf{k}_i$ to indicate the transverse momentum vectors of the gluons coupled to the photon.

$$D = (\epsilon^2 + \mathbf{k}^2)(\epsilon_1^2 + (\mathbf{k} + \mathbf{k}_1 - \alpha \mathbf{l})^2) \quad (\text{A.35})$$

$$\epsilon^2 = Q^2 \alpha (1 - \alpha) + m_f^2, \quad \epsilon_1^2 = (Q^2 + l^2) \alpha (1 - \alpha) + m_f^2 \quad (\text{A.36})$$

e_f and m_f are the quark electric charge and mass and the numerator for a transversal photon is

$$N^\perp = m_f^2 + (\alpha^2 + (1 - \alpha)^2)\mathbf{k}(\mathbf{k} + \mathbf{k}_1) - \alpha^2\mathbf{k}\mathbf{l} \quad (\text{A.37})$$

and for a longitudinal photon is

$$N^L = 4Q^2\alpha^2(1 - \alpha)^2 \quad (\text{A.38})$$

This expression can be conveniently represented as an integral over the colour dipole density ρ created by the $q\bar{q}$ pair at a given distance in the transverse space:

$$f(k_1, k_2) = \int d^2r \rho_l(r) e^{ik_1 r} \quad (\text{A.39})$$

From (A.34) - (A.38) one finds for the transverse and longitudinal photons:

$$\begin{aligned} \rho_l^\perp(r) &= \frac{e_f^2}{(2\pi)^3} e^{-i\alpha l r} \int_0^1 d\alpha \left[m_f^2 K_0(\epsilon r) K_0(\epsilon_1 r) + (\alpha^2 + (1 - \alpha)^2) \epsilon \epsilon_1 K_1(\epsilon r) K_1(\epsilon_1 r) - \right. \\ &\quad \left. \alpha(1 - \alpha)(1 - 2\alpha) \frac{i\epsilon \mathbf{l} \mathbf{r}}{r} K_0(\epsilon_1 r) K_1(\epsilon r) \right] \\ \rho_l^L(r) &= \frac{4e_f^2 Q^2}{(2\pi)^3} e^{-i\alpha l r} \int_0^1 d\alpha \alpha^2 (1 - \alpha)^2 K_0(\epsilon r) K_0(\epsilon_1 r) \end{aligned} \quad (\text{A.40})$$

A.3 Coordinate representation of the non-forward G function

For the non-forward direction a function $G(k_1, k_2, k_3)$ can be defined by a natural generalization of the infrared stable function introduced by J.Bartels [20]:

$$G(k_1, k_2, k_3) = -g^2 N_c W_2(k_1, k_2, k_3) - D(k_1, k_2 + k_3)(\omega(k_2) - \omega(k_2 + k_3)) - D(k_1 + k_2, k_3)(\omega(k_2) - \omega(k_1 + k_2)) \quad (\text{A.41})$$

where $D(k_1, k_2)$ is the amputated BFKL function with the gluon momenta k_1 and k_2 ; $\omega(k)$ is the gluon trajectory and $W_2(k_1, k_2, k_3)$ is the $2 \rightarrow 3$ vertex integrated with the function D . A graphical representation of the function G , following the notation introduced by Bartels, has been given in Fig. 3.1. G can be presented as a sum of proper (integral) G_1 and improper (terms with ω) G_2 parts which, by definition, are given by

$$G_2(k_1, k_2, k_3) = -D(k_1, k_2 + k_3)(\omega(k_2) - \omega(k_2 + k_3)) - D(k_1 + k_2, k_3)(\omega(k_2) - \omega(k_1 + k_2)) \quad (\text{A.42})$$

and

$$G_1(k_1, k_2, k_3) = g^2 N_c \int \frac{d^2 q_1 d^2 q_3}{(2\pi)^3} \delta^2(q_1 + q_3 - k_1 - k_2 - k_3) D(q_1, q_3) \left(\frac{(k_2 + k_3)^2}{(q_1 - k_1)^2 q_3^2} + \frac{(k_1 + k_2)^2}{q_1^2 (q_3 - k_3)^2} - \frac{k_2^2}{(q_1 - k_1)^2 (q_3 - k_3)^2} - \frac{(k_1 + k_2 + k_3)^2}{q_1^2 q_3^2} \right) \quad (\text{A.43})$$

For the following we recall that the gluon trajectory is

$$\omega(k) = -(1/2)g^2 N_c \int \frac{d^2 q}{(2\pi)^3} \frac{k^2}{q^2 (q - k)^2} = -\frac{g^2 N_c}{8\pi^2} \ln \frac{k^2}{m^2} \quad (\text{A.44})$$

where the gluon mass m is a regularization parameter (to be set to zero in the infrared finite expressions).

We shall consider the Fourier transform

$$G(r_1, r_2, r_3) = \frac{1}{(2\pi)^6} \int d^2 k_1 d^2 k_2 d^2 k_3 \exp(i \sum_{j=1}^3 k_j r_j) G(k_1, k_2, k_3) \quad (\text{A.45})$$

To simplify the notation we denote

$$d\tau(k) = \frac{1}{(2\pi)^6} d^2 k_1 d^2 k_2 d^2 k_3 \quad , \quad d\tau(q) = \frac{d^2 q_1 d^2 q_3}{(2\pi)^3} \delta^2(q_1 + q_3 - k_1 - k_2 - k_3) \quad (\text{A.46})$$

and

$$d\tau(\rho) = d^2 \rho_1 d^2 \rho_3 \quad (\text{A.47})$$

A.3.1 Proper part

Let us start by considering the terms (four) which appear in the proper part (A.43) and which will be denoted as G_{11} , G_{12} , ... etc. For the first term one has

$$G_{11} = g^2 N_c \int d\tau(k) d\tau(q) d\tau(\rho) D(\rho_1, \rho_3) \frac{(k_2 + k_3)^2}{(q_1 - k_1)^2 q_3^2} \exp(i \sum_{j=1}^3 k_j r_j - i q_1 \rho_1 - i q_3 \rho_3) \quad (\text{A.48})$$

After the substitution $(k_2 + k_3)^2 \rightarrow -(\partial_2 + \partial_3)^2$, $1/q_3^2 \rightarrow -\partial_{\rho_3}^{-2}$ and $q_1 - k_1 \rightarrow k_2$ (all operators ∂ refer to coordinate space), choosing q_3 as independent variable one obtains

$$G_{11} = g^2 N_c (\partial_2 + \partial_3)^2 \int d\tau(k) d\tau(q) d\tau(\rho) \frac{1}{k_2^2} \exp(ik_1 r_1 + ik_3 r_3 + i(q_3 - k_3 + k_2)r_2 - i(k_1 + k_2)\rho_1 - iq_3 \rho_3) \partial_{\rho_3}^{-2} D(\rho_1, \rho_3) \quad (\text{A.49})$$

Integrations over k_1 , k_3 and q_3 give $(2\pi)^2 \delta^2(r_1 - \rho_1)$, $(2\pi)^2 \delta^2(r_2 - r_3)$ and $(2\pi)^2 \delta^2(r_2 - \rho_3)$ respectively and the remaining integral over k_2 has to be regularized:

$$\int \frac{d^2 k_2}{k_2^2 + m^2} \exp i k_2 (r_2 - \rho_1) = 2\pi (c - \ln |r_2 - \rho_1|), \quad m \rightarrow 0, \quad (\text{A.50})$$

where

$$c = \ln(2/m) + \psi(1) \quad (\text{A.51})$$

Doing the integrations by means of δ -functions, we obtain

$$G_{11} = \frac{g^2 N_c}{4\pi^2} (\partial_2 + \partial_3)^2 \delta^2(r_{23}) (c - \ln r_{12}) \partial_3^{-2} D(r_1, r_3) \quad (\text{A.52})$$

Here and in the following $r_{ij} = r_i - r_j$. Note that the operator $\partial_2 + \partial_3$ does not act on the δ -function. Its action on the $\ln r_{12}$ and D is evidently reduced to ∂_2 and ∂_3 respectively. However in the end one has to put $r_2 = r_3$ so that we obtain finally

$$G_{11} = \frac{g^2 N_c}{4\pi^2} \delta^2(r_{23}) \partial_3^2 (c - \ln r_{13}) \partial_3^{-2} D(r_1, r_3) \quad (\text{A.53})$$

The Fourier transform of the second term in (A.43) is obtained from G_{11} by interchanging $1 \leftrightarrow 3$:

$$G_{12} = \frac{g^2 N_c}{4\pi^2} \delta^2(r_{12}) \partial_1^2 (c - \ln r_{13}) \partial_1^{-2} D(r_1, r_3) \quad (\text{A.54})$$

Now we consider to the Fourier transform of the third term in (A.43). Its explicit form is

$$G_{13} = -g^2 N_c \int d\tau(k) d\tau(q) d\tau(\rho) D(\rho_1, \rho_3) \frac{k_2^2}{(q_1 - k_1)^2 (q_3 - k_3)^2} \exp(i \sum_{j=1}^3 k_j r_j - i q_1 \rho_1 - i q_3 \rho_3) \quad (\text{A.55})$$

As usual we substitute $k_2^2 \rightarrow -\partial_2^2$, $k_1 \rightarrow k_1 + q_1$, $k_3 \rightarrow k_3 + q_3$ and choose k_2 as a dependent

variable, obtaining

$$G_{13} = g^2 N_c \partial_2^2 \int d\tau(k) d\tau(q) d\tau(\rho) D(\rho_1, \rho_3) \frac{1}{k_1^2 k_3^2} \exp(i(k_1 + q_1)r_1 + i(k_3 + q_3)r_3 - i(k_1 + k_3)r_2 - iq_1\rho_1 - iq_3\rho_3) \quad (\text{A.56})$$

The integrals over q_1 and q_3 give $(2\pi)^2 \delta^2(r_1 - \rho_1)$ and $(2\pi)^2 \delta^2(r_3 - \rho_3)$ respectively. The two integrals over k_1 and k_3 give $2\pi(c - \ln r_{12})$ and $2\pi(c - \ln r_{23})$, so that one has

$$G_{13} = \frac{g^2 N_c}{8\pi^3} \partial_2^2 (c - \ln r_{12})(c - \ln r_{23}) D(r_1, r_3) \quad (\text{A.57})$$

and, after performing the derivatives in r_2 ,

$$G_{13} = -\frac{g^2 N_c}{8\pi^3} \left(2 \frac{\mathbf{r}_{12} \mathbf{r}_{23}}{r_{12}^2 r_{23}^2} + 2\pi(c - \ln r_{13})(\delta^2(r_{23}) + \delta^2(r_{12})) \right) D(r_1, r_3) \quad (\text{A.58})$$

We are left with the last proper term

$$G_{14} = -g^2 N_c \int d\tau(k) d\tau(q) d\tau(\rho) D(\rho_1, \rho_3) \frac{(k_1 + k_2 + k_3)^2}{q_1^2 q_3^2} \exp(i \sum_{j=1}^3 k_j r_j - iq_1 \rho_1 - iq_3 \rho_3) \quad (\text{A.59})$$

Substituting all k 's by the derivatives w.r.t. r 's and q 's by the derivatives w.r.t. ρ 's we can do all the integrals to obtaining

$$G_{14} = \frac{g^2 N_c}{2\pi} (\partial_1 + \partial_2 + \partial_3)^2 \delta^2(r_{12}) \delta^2(r_{23}) \partial_1^{-2} \partial_3^{-2} D(r_1, r_3) \quad (\text{A.60})$$

Since the sum $\sum_{i=1}^3 \partial_i$ does not act on the two δ - functions, we can transfer the first operator to the right and note that, acting on D , ∂_2 gives zero. So finally one has

$$G_{14} = \frac{g^2 N_c}{2\pi} \delta^2(r_{12}) \delta^2(r_{23}) (\partial_1 + \partial_3)^2 \partial_1^{-2} \partial_3^{-2} D(r_1, r_3) \quad (\text{A.61})$$

A.3.2 Improper part

A similar notation is used for the improper part: $G_{21}, G_{22}, \dots$ correspond to the 4 terms in (A.42). Take the first term

$$G_{21} = - \int d\tau(k) d\tau(\rho) \omega(k_2) D(\rho_1, \rho_3) \exp(i \sum_{j=1}^3 k_j r_j - ik_1 \rho_1 - i(k_2 + k_3) \rho_3) \quad (\text{A.62})$$

We transform this expression in two possible ways. One is evident: we just take ω out of the integral as an operator to obtain immediately

$$G_{21} = -\omega(-i\partial_2) \delta^2(r_{23}) D(r_1, r_3) \quad (\text{A.63})$$

In this form it is clear that the contribution can be split into a sum of two independent part, a holomorphic one acting on r 's and an antiholomorphic one acting on r^* 's. However this expression is difficult to analyze as to its conformal properties, due to the fact that the operator ω is applied to the δ -function, that is, to both D and a function which is to be integrated with G

A convenient alternative form for G_{21} is obtained by means of integration over k_1 and k_3 in (A.62)

$$G_{21} = -D(r_1, r_3) \int \frac{d^2 k_2}{(2\pi)^2} e^{ik_2 r_{23}} \omega(k_2) = -\tilde{\omega}(r_{23}) D(r_1, r_3) \quad (\text{A.64})$$

where $\tilde{\omega}(r)$ is just the Fourier transform of $\omega(k)$.

A useful relation, similar to (A.50),

$$\int \frac{d^2 r}{r^2 + \epsilon^2} e^{ikr} = 2\pi(c_1 - \ln k) \quad , \quad c_1 = \ln(2/\epsilon) + \psi(1) \quad (\text{A.65})$$

allows us to write

$$\omega(k) = -\frac{g^2 N_c}{8\pi^2} \ln \frac{k^2}{m^2} = \frac{g^2 N_c}{8\pi^3} \left[\int \frac{d^2 r}{r^2 + \epsilon^2} e^{ikr} - 2\pi(c_1 - \ln m) \right] \quad (\text{A.66})$$

On noting that

$$c_1 - \ln m = c - \ln \epsilon \quad (\text{A.67})$$

from eq. (A.66) we conclude that

$$\tilde{\omega}(r) = \frac{g^2 N_c}{8\pi^3} \left[\frac{1}{r^2 + \epsilon^2} + 2\pi(\ln \epsilon - c)\delta^2(r) \right] \quad (\text{A.68})$$

It also follows that the right-hand side of (A.68) does not, in fact, depend on ϵ . Evidently the term with $\ln \epsilon$ serves to regularize the singularity of the first term at $r = 0$. In the following this regularization will always be implied, so that

$$\frac{1}{r^2} \equiv \frac{1}{r^2 + \epsilon^2} + 2\pi \ln \epsilon \delta^2(r), \quad \epsilon \rightarrow 0, \quad (\text{A.69})$$

With this convention we finally obtain

$$\tilde{\omega}(r) = \frac{g^2 N_c}{8\pi^3} \left(\frac{1}{r^2} - 2\pi c \delta^2(r) \right) \quad (\text{A.70})$$

The same result could be obtained directly by Fourier transforming the integral representation of $\omega(k)$ but in this way we obtain (A.69) as a byproduct. Using this we get our final expression

$$G_{21} = -\frac{g^2 N_c}{8\pi^3} \left(\frac{1}{r_{23}^2} - 2\pi c \delta^2(r_{23}) \right) D(r_1, r_3) \quad (\text{A.71})$$

The term G_{23} is obtained from this by the permutation 1 and 3 in the operator acting on D :

$$G_{23} = -\frac{g^2 N_c}{8\pi^3} \left(\frac{1}{r_{12}^2} - 2\pi c \delta^2(r_{12}) \right) D(r_1, r_3) \quad (\text{A.72})$$

Improper terms 2 and 4 are calculated in a simpler manner. In fact we have

$$G_{22} = - \int d\tau(k) d\tau(\rho) \omega(k_2 + k_3) D(\rho_1, \rho_3) \exp(i \sum_{j=1}^3 k_j r_j - i k_1 \rho_1 - i(k_2 + k_3) \rho_3) \quad (\text{A.73})$$

In this case we can substitute $\omega(k_2 + k_3)$ by an operator $\omega(-i\partial_{\rho_3})$ acting on D . After that all integrations are trivially done and we obtain

$$G_{22} = \delta^2(r_{23}) \omega(-i\partial_3) D(r_1, r_3) \quad (\text{A.74})$$

and in a similar manner

$$G_{24} = \delta^2(r_{12}) \omega(-i\partial_1) D(r_1, r_3) \quad (\text{A.75})$$

A.3.3 Infrared stable pieces

Let us see how to combine different proper and improper terms in order to obtain expressions which do not depend on m and so are infrared finite. There are three infrared finite combinations.

Summing G_{11} and G_{22} we find

$$G^{(1)} = G_{11} + G_{22} = \delta^2(r_{23}) \frac{g^2 N_c}{8\pi^2} \left[2 \ln 2 + 2\psi(1) - k_3^2 \ln r_{13}^2 k_3^{-2} - \ln k_3^2 \right] D(r_1, r_3) \quad (\text{A.76})$$

Here we have used the explicit expression for ω in the momentum space and also used the notation k_3 for the operator $-i\partial_3$ acting in coordinate space. The important point is that the terms with $\ln m$ have cancelled in the sum, so that the expression is infrared finite. Note that the expression in the brackets is a half of the Hamiltonian for the two gluons (with a minus sign). It represents the Hamiltonian of the gluon 3 in interaction with the gluon 1 considered as an external source.

The other half of the Hamiltonian appears in the sum G_{12} and G_{24} , for which we also find an infrared finite expression

$$G^{(2)} = G_{12} + G_{24} = \delta^2(r_{12}) \frac{g^2 N_c}{8\pi^2} \left[2 \ln 2 + 2\psi(1) - k_1^2 \ln r_{13}^2 k_1^{-2} - \ln k_1^2 \right] D(r_1, r_3) \quad (\text{A.77})$$

The rest of the terms give an expression

$$\begin{aligned} G^{(3)} &= G_{13} + G_{14} + G_{21} + G_{23} \\ &= \frac{g^2 N_c}{8\pi^3} \left[-4\pi^2 \delta^2(r_{12}) \delta^2(r_{23}) (k_1 + k_3)^2 k_1^{-2} k_3^{-2} + 2\pi (\delta^2(r_{23}) + \delta^2(r_{12}) \ln r_{13} - \frac{r_{13}^2}{r_{12}^2 r_{23}^2}) \right] D(r_1, r_3) \end{aligned} \quad (\text{A.78})$$

It is also infrared finite since all the $\ln m$ terms cancel. One has only to remember that the

singularities of the last term at $r_{12} = 0$ and $r_{23} = 0$ have to be regularized, if necessary, according to the prescription (A.69).

A.4 Some useful relations in graphical notation.

We introduce in the following a graphical notation in order to easily derive some relations involving the elementary reggeized gluons (see also in [29] for the analytical counterpart). A line will denote a reggeized gluon momentum, a line split a dependence in the sum of the momenta corresponding to the lines, the squares the interaction vertices defined in (3.2)-(3.4) and a small circle the contribution of a Regge trajectory (if more than one is present a sum of all the possible one-circle graphs is meant). In Fig. A.1 we show the vertices with their explicit definition, related to the $2 \rightarrow 2$, $2 \rightarrow 3$ and $2 \rightarrow 4$ transitions in (a), (b) and (c) respectively, and in Fig. A.2 a possible representation of the bootstrap condition.

It is simple to check the validity of the relation (3.33) which was needed to calculate the triple pomeron vertex Z . The steps are shown in Fig. A.3 and depend exclusively on the repeated application of the relation (b) of Fig. A.1. The relation in line (a) is directly derived by applying on $W(4, 1, 3)$ (first term in the rhs) this relation with the indices $(123) \rightarrow (413)$. In (b) and (c) in a similar manner other useful relations are derived and, applying them to (a), one finds in (d) the identity (3.33).

Let us show in this graphical representation how to apply the bootstrap condition, giving the proof that the solution of eq. (3.11) is in fact the expression (3.13). The proof is based on the reduction, by means of the bootstrap, of the equation for the 3 gluon system to the equation for two gluon in the singlet state, i.e. a BFKL non-homogeneous equation. Precisely one is able to show this for an equation with an inhomogeneous term which is a part of the one present in (3.11) so one completes the solution by applying the linearity of the equation. Therefore we decompose the second inhomogeneous term $D_{2 \rightarrow 3}$ in a sum of three terms as in D_{30} and study one of the three similar equations, corresponding to the case with $D_{20}(1)$ and the third and fourth terms in the decomposition of $D_{2 \rightarrow 3}$ given in Fig. A.4.a.

We can now easily check that the solution of the resulting equation (Fig. A.4.b) is given in fact by $D_3^{(1)} = -D_2(1, 2 + 3)$. With this substitution the equation is represented in Fig. A.4.c for the D_2 . On applying the bootstrap condition one has the relation in Fig. A.4.d and in Fig. A.4.e two relations are derived from Fig. A.1.b. Substituting these identities in the relation of Fig. A.4.c one finally obtains for D_2 the equation in Fig. A.4.f which is the expected BFKL equation.




Figure A.1: Graphical representation of the elementary vertices.







Figure A.2: Graphical representation of the bootstrap condition.

(a) 

(b) 

(c) 






(d) 


Figure A.3: Proof of the relation (3.33).


(a) 

(b) $(j-1)$ 

(c) $(j-1)$ 

(d) 

(e) 




(f) $(j-1)$ 

Figure A.4: Proof of the relation (3.13).

Bibliography

- [1] G. Veneziano, Phys. Lett. **B52** (1974) 220; Nucl. Phys. **B117** (1976) 519.
- [2] G. t'Hooft, Nucl. Phys. **B72** (1979) 461.
- [3] E. Witten, Nucl. Phys. **B160** (1979) 57.
- [4] L.N.Lipatov, Yad. Fiz. **23** (1976) 642.
- [5] V.S.Fadin, E.A.Kuraev and L.N.Lipatov, Phys. Lett. **B60** (1975)50;
I.I.Balitsky and L.N.Lipatov, Sov.J.Nucl.Phys. **15** (1978) 438.
- [6] L.N. Lipatov and V.S. Fadin, Sov. J. Nucl. Phys. **50** (1989) 712
- [7] V.S. Fadin, R. Fiore and M.I. Kotsky, Phys. Lett. **B359** (1995) 181, **B387** (1996) 593;
V.S. Fadin and L.N. Lipatov, Nucl. Phys. **B406** (1993) 259, Nucl. Phys. **B477** (1996) 767;
V.S. Fadin, M.I. Kotsky, L.N. Lipatov, Phys. Lett. **B415** (1997) 97.
- [8] V.S. Fadin and L.N. Lipatov, preprint hep-ph**9802290**.
- [9] I. Abt et al., H1 Collab.,Nucl. Phys. **B407** (1993) 515;
T. Ahmed et al., H1 Collab., DESY 95-006 (1995);
M. Derrick et al., ZEUS Collab., Phys. Lett. **B316** (1993) 412;
M. Derrick et al., ZEUS Collab., DESY 94-143 (1994).
- [10] ZEUS Collaboration, PREPRINT, Measurement of the F_2 structure function in deep in-elastic e^+p scattering using 1994 data from ZEUS detector at HERA.
- [11] A.H. Mueller, Nucl.Phys. **B415** (1994) 373; **B437** (1995) 107.
- [12] A.H. Mueller and B.Patel, Nucl. Phys. **B425** (1994) 471.
- [13] M.A.Braun, Phys. Lett. **B348** (1995) 190.
- [14] M.A.Braun, Phys. Lett. **B345** (1995) 155.
- [15] M.A.Braun, G.P. Vacca and G. Venturi, Phys. Lett. **B388** (1996) 823.
- [16] M.A. Braun, G.P. Vacca, Phys. Lett. **B406** (1997) 137-148.
- [17] M.A. Braun, G.P. Vacca, EPJ **C ?** (1998), preprint hep-ph/ **9706314**.

- [18] L.N. Gribov, E.M. Levin and M.G. Ryskin, Phys. Rep. **100** (1983) 1.
- [19] J. Bartels, Z. Phys. **C60** (1993) 471.
- [20] J.Bartels and M.Wuesthoff, Z.Phys. **C 66** (1995) 157.
- [21] J.Bartels, L.N.Lipatov and M.Wuesthoff, Nucl. Phys. **B464** (1996) 298.
- [22] J.Bartels, Nucl. Phys. **B151** (1979) 293.
- [23] J. Bartels, Nucl. Phys. **B175** (1980) 365.
- [24] L.N. Lipatov, Phys. Lett. **B309** (1993) 394; JETP Lett **59** (1994) 596.
- [25] L.D. Faddeev and G.P. Korchemsky, Phys. Lett. **B342** (1995) 311.
- [26] V.O. Tarasov, L.A. Takhtajan, L.D. Faddeev, Theor. Math. Phys. **57** (1983) 163.
- [27] L.N.Lipatov, in "Perturbative QCD", ed. A.H.Mueller (World Sci., Singapore, 1989).
- [28] G.P. Korchemsky, preprint LPTHE-Orsay-97-62, hep-ph/9711277.
- [29] M.A.Braun, S.Petersburg preprint SPbU-97-9 (hep-ph/9706373).
- [30] M.A. Braun and G.P. Vacca, Bologna Univ. preprint, hep-ph/9711486.
- [31] R.Peschanski, Saclay preprint, (hep-ph/9704342); Phys.Lett. **B409** (1997) 491.
- [32] H. Cheng, C. Y. Lo, Phys. Rev. **D15** (1977) 2959.
- [33] M.Grisaru, H.Schnitzer and H.S.Tsao, Phys. Rev.,**D12** (1973) 4498.
- [34] P. Higgs, Phys. Lett. **12** (1964) 132; Phys. Rev. Lett. **13** (1964) 508.
- [35] F. Englert and R. Brout, Phys. Rev. Lett. **13** (1964) 3.
- [36] J.C. Collins, D.E. Soper and G. Sterman, in "Perturbative QCD", ed. A.H.Mueller (World Sci., Singapore, 1989).
- [37] V.N. Gribov and L.N. Lipatov, Sov. J. nucl. Phys., **15** (1972) 438,675;
G. Altarelli and G. Parisi, Nucl. Phys. **B126** (1977) 297;
Y. Dokshitzer, Sov. Phys. JETP **46** (1977) 641.
- [38] S. Catani and F. Hautmann, Nucl. Phys. **B427** (1994) 475;
S. catani, M. Ciafaloni and F. Hautmann, Phys. Lett **B242** (1990) 97; Nucl. Phys. **B 366**
(1991) 135.
- [39] M.A. Braun , Santiago preprint, *The interaction of reggeized gluons and Lipatov's hard pomeron* (1994).
- [40] L.N. Lipatov, Sov. Phys. JETP **63** (1986) 904.

- [41] M. Froissart, Phys. Rev. **12** (1961) 1053
- [42] W. Butchmüller, D. Haidt, hep-ph/9605428
- [43] A.L. Ayala, M.B. Gay Ducati, E.M. Levin, Phys. Lett **B388** (1996) 188.
- [44] N.Nikolaev and B.Zakharov, Phys.Lett **B327** (1994) 157.
- [45] E.M. Levin, M.G. Ryskyn, Phys. Rep. **189**, **N.6** (1990), 267-382.
- [46] N.N.Nikolaev and B.G.Zakharov, Z.Phys. **C49** (1991) 607.
- [47] R.E. Handcock, D.A. Ross, Nucl. Phys. **383** (1992) 575
- [48] A.J.Askew, J. Kwiecinski, A.D. Martin, P.J.Sutton, Phys. Rev. **D47** (1993) 3775, Phys. Rev. **D49** (1994) 4402;
A.J.Askew, K. Golec-Biernat, J. Kwiecinski, A.D. Martin, P.J.Sutton, Phys. Lett. **B325** (1994) 212.
- [49] G.P. Vacca and G. Venturi, Il Nuovo Cimento **108A**, (1995) 77-85.
- [50] M.A.Braun, Phys. Lett. **B357** (1995) 138.
- [51] L.V. Gribov, E.M. Levin, M.G. Ryskyn, Phys. Lett. **B121** (1983) 65-71.
- [52] H1 Collab., Nucl. Phys. **B439** (1995) 471.
- [53] See M.Block et al in *Proc. 24th Int. Symp. on Multiparticle Dynamics*, Ed. A.Giovanini, World Scient. (1995).
- [54] M. Glück, E. Reya, A. Vogt, Z. Phys **C67** (1995) 433-447.
- [55] A.D. Martin, W.J. Stirling, R.G. Roberts, Phys. Rew. **D50** (1994) 6734.
- [56] J. Kwiecinski, A.D. Martin, P.J. Sutton, Phys. Rew. **D44** (1991) 2640.
- [57] H. Lotter, PhD Thesis, hep-ph/9705288.
- [58] A. Bialas, H. Navelet and R. Peschanski, Saclay preprints hep-ph/9711236 and hep-ph/9711442.

UNIVERSITÀ DEGLI STUDI DI BOLOGNA

FACOLTÀ DI SCIENZE MATEMATICHE, FISICHE E NATURALI

Dottorato di Ricerca in Fisica

The hard QCD pomeron:
some aspects of its phenomenology and interactions

Tesi di dottorato di:

Gian Paolo Vacca

Coordinatore:

Chiar.mo Prof.
Giulio Pozzi

Supervisore:

Chiar.mo Prof.
Giovanni Venturi

X ciclo (1994-1997)